

## Accommodation to uncomfortable patterns

Authors: Sarah M. Haigh<sup>1</sup>, Wolfgang Jaschinski<sup>2</sup>, Peter M. Allen<sup>3</sup>, & Arnold J. Wilkins<sup>4</sup>

<sup>1</sup> Visual Perception Unit, Department of Psychology, University of Essex, Wivenhoe Park, Colchester, Essex, CO 4 3SQ, UK. [shaigh@essex.ac.uk](mailto:shaigh@essex.ac.uk).

<sup>2</sup> Leibniz-Institute für Arbeitsforschung an der TU Dortmund, Ardeystr. 67, D-44139, Dortmund, Germany. [jaschinski@ifado.de](mailto:jaschinski@ifado.de).

<sup>3</sup> Vision and Eye Research Unit, Postgraduate Medical Institute, Anglia Ruskin University, Cambridge, CB1 1PT, UK. [peter.allen@anglia.ac.uk](mailto:peter.allen@anglia.ac.uk).

<sup>4</sup> Visual Perception Unit, Department of Psychology, University of Essex, Wivenhoe Park, Colchester, Essex, CO 4 3SQ, UK. [arnold@essex.ac.uk](mailto:arnold@essex.ac.uk).

Keywords: accommodation, pattern sensitivity, gratings

Address for correspondence

Arnold J Wilkins,  
Visual Perception Unit,  
Department of Psychology,  
University of Essex,  
Colchester, CO4 3SQ.

## Abstract

Grating patterns can cause discomfort and perceptual distortions. Individuals who experience discomfort and are susceptible to these distortions generally show weaker accommodation than those who are less susceptible. We measured the accommodative response to grating patterns known to differ in the discomfort they evoke because of differences in their colour, motion or spatial frequency. The parameters known to affect discomfort and distortion had no influence on the mean or variance in the accommodative response, even when accommodative demand was manipulated systematically and the accommodative response varied as expected.

## 1. Introduction

In patients with photosensitive epilepsy, patterns of stripes can evoke paroxysmal epileptiform electroencephalographic activity. The patterns to which patients are susceptible are high in contrast and have a spatial frequency between 0.5 and 12 cycles/degree (cpd) with sensitivity maximal at about 3cpd. Such patterns are generally perceived as uncomfortable to view by healthy individuals and can evoke visual illusions. Some individuals are more susceptible to the illusions than others, and these individuals are generally more susceptible to headaches (Wilkins *et al.*, 1984). Illusions are sometimes reported when reading text, and the illusions seen in text can resemble those seen in gratings: a rhomboid lattice, rainbow colours, and perceptual instability (Wilkins and Nimmo-Smith, 1987). Some aspects of the perceptual instability have been explained as due to accommodative fluctuations (Helmholtz, 1856; Millodot, 1968; Campbell & Robson, 1958).

Accommodative fluctuations have also been proposed as an explanation for a variety of visual illusions, including the distortions seen in fine grating patterns (Helmholtz, 1856, 1924), concentric circles (Helmholtz, 1856, 1924; Purkinje, 1823, 1825), and in the 'fluttering hearts' illusion. Purkinje (1823, 1825) argued that the distortions (specifically in concentric circles) were due to an unfocused image on the retina, but it was Helmholtz (1924) who specifically identified the accommodative system as being potentially responsible for the blurred and distorted image. Brewster, Wheatstone and Helmholtz (Wade, 1983) observed that a red shape (in this case a heart shape) on a green background created an illusion of depth in which the background and the shape appeared to oscillate, causing a 'fluttering' illusion. There has been much speculation as to why this illusion might occur. One possibility is that owing to longitudinal chromatic aberration of the eye the normal fluctuations in accommodation become detectable. Another possibility is that the accommodative fluctuations are exaggerated because the accommodative system is "colour blind" (Wolfe & Owens, 1981) and cannot detect the contours of the shape from the background, particularly when the contour is isoluminant. Isoluminance has an effect on form perception (Gregory, 1977), despite the isoluminant chromatic contours being well above detection threshold (Switkes, Bradley & Schor, 1990).

In the majority of studies that have measured the accommodative response to isoluminant patterns the variance in the response has not been reported. Wolfe and Owen (1981) found that the accommodative response to isoluminant contours was much weaker compared to contours with high luminance contrast. They did not record accommodation continuously, but remarked that they did not see any great fluctuations in the accommodative response over time. Switkes, Bradley and

**Deleted:** in the context of the 'fluttering hearts' illusion.

Schor (1990) similarly found a poor accommodative response to isoluminant contours compared to contours with high luminance contrast, but did not report the variance in the accommodative response (even though they recorded continuously at 20Hz).

The difference between the accommodative response and accommodative stimulus appropriate for the actual viewing distance is known as the accommodative lag and varies from one individual to another: the unit of these measures is the dioptre, i.e. the inverse measure of focal distance in metres. Individuals with large accommodative lags tend to report discomfort and illusions. Allen *et al.* (2010) measured the effect of an achromatic 1.3cpd grating pattern on accommodative lag in participants who reported seeing a large number of illusions in a pattern of stripes. They found that these participants had a larger accommodative lag (they under-accommodated for the viewing distance) than those who reported few illusions. When the participants used a tinted overlay of their choice, the lag and the discomfort were both reduced. Simmers, Gray and Wilkins (2001) also found that tinted lenses reduced the variability in the accommodative response in those who reported discomfort and illusions. Therefore, it is possible that it is the weaker accommodative response (and the consequent blur) that causes the illusions and discomfort.

Similarly, Chase *et al.* (2009) found a positive correlation between the accommodative lag in response to a 0.2m target and the visual discomfort scores from the Conlon Visual Discomfort Survey. Tosha *et al.* (2009) also found that a group which reported high discomfort showed greater fatigue when sustaining accommodation at near distance (3-4D), even over a short time period (90s).

A poor accommodative response does not appear to explain all of the illusions that have previously been reported in stripes. A rhomboid lattice illusion is often seen (Wilkins, 1986). The size of the lattice appears to be affected by the spatial frequency of the target pattern (Wilkins & Nimmo-Smith, 1987). It cannot readily be explained by peripheral factors and an explanation in terms of cortical inhibition has been proposed by Georgeson (1976, 1980).

An alternative theory for the cause of the visual discomfort and illusions seen in striped patterns is that the cortex produces a heightened neural response to the pattern (Huang *et al.*, 2003). It is possible that the excess firing can spread locally to nearby parts of the cortex and can cause the appearance of illusions. Adjajian *et al.* (2004) found maximal cortical power in the gamma frequencies to patterns which caused the most discomfort and illusions. Siniatchkin *et al.* (2007) also found larger N75-P100 and P100-P135 components in the visual evoked potential in response to uncomfortable patterns in those individuals with a propagating photoparoxysmal response but not in those with local photoparoxysmal response. This could explain why some individuals are more

susceptible to illusions in grating patterns: those with a hyperexcitable cortex (e.g. individuals with migraine) produce an over-response to the patterns causing more illusions.

In the following four experiments, we investigated the differences in the accommodative response to gratings that differed with respect to pattern parameters known to affect how uncomfortable the patterns are to view. The first three studies investigated the accommodative lag in response to grating patterns that varied in parameters of colour or movement. In the final study, we varied the accommodative demand, and investigated the differences in accommodative lag in response to grating patterns that varied in their spatial frequency and contrast. If a poor accommodative response is responsible for the discomfort, then one might anticipate an association between the patterns that normally evoke discomfort and the accommodative lag when viewing these patterns.

## 2.1 Experiment 1 Accommodation to coloured gratings

Chromatic contrast, as well as luminance contrast, is known to contribute to visual discomfort (Haigh et al., 2012a; Haigh et al., 2012b; Wilkins et al., 2008). Wilkins et al. (2008) measured the ratings of discomfort to a series of chromatic grating patterns that varied in the separation in chromaticity of the component bars. Participants were asked to rate how uncomfortable the pattern was to view on a Likert scale. The gratings varied in the hue and saturation of the component bars and in all experiments, there was a linear increase in aversion with the separation in CIE UCS chromaticity irrespective of luminance contrast and of the hue of the bars. In principle, it is possible that the large separations in chromaticity could be causing a weak accommodative response and/or large fluctuations in accommodation because of longitudinal chromatic aberration: the accommodative system might attempt to accommodate to two different distributions of spectral power.

In the following experiment, the mean chromaticity of a grating was varied systematically. Three types of grating were presented, each having colours formed from just two of the three colour pixels of an LCD display: i.e. with average chromaticities on the yellow, turquoise and purple lines at the extremes of the display gamut. The accommodative response was measured using an open-field autorefractor.

## 2.2 Method

### 2.2.1 Participants

All studies were approved by the University of Essex Ethics Committee, Essex, UK, and adhere to the tenets of the Declaration of Helsinki (Oct, 2008). Four male and 26 female undergraduate students

**Deleted:** In six experiments, Haigh, Tang, Coutts and Wilkins (submitted)


aged 18-33 (mean 20) from the University of Essex took part. All had a minimum acuity of 6/6 monocularly at near and binocularly at distance and near (Lighthouse Near and Far tests of visual acuity) and a minimum stereoacuity of 60sec.arc (Titmus test). Log contrast sensitivity for letters was at least 2.00 (Pelli-Robson letter chart), and no colour deficiencies were detected (Ishihara plates). None of the participants required prescription glasses or contact lenses.

Participants completed a questionnaire about their general health, visual history and the headaches they experienced. None of the participants had a history of seizures.

### 2.2.2 Stimuli

A telespectroradiometer (model PR-670®, Photo Research®, Chatsworth, CA, USA) was used to measure the ( $u'$   $v'$ ) chromaticity of the red, blue and green pixels from a Dell Precision M4500 laptop screen. The chromaticities used for the gratings were made up of only two of the red, blue or green pixels. The chromaticity of the point mid-way between the chromaticities of the red pixel and the green pixel was found. Pairs of colours (with the same photometric luminance, 23 cd.m<sup>-2</sup>) that lay either side of the mid-point and equidistant from it were used in alternating bars in a 2cpd square-wave grating pattern. The colour distance between the paired chromaticities was gradually increased to make seven gratings. For example, the smallest colour difference was between two shades of yellow. The colour difference increased to make a grating with bars of yellow-red and yellow-green, with the largest colour difference being a grating with bars of a red and a green. This was repeated for the red-blue and the blue-green colour pairs to create a total of 21 gratings. The chromaticities of the colours used are shown in Figure 1.

**Deleted:** All bars had similar luminance (23cd.m<sup>2</sup>).



**Figure 1.** CIE UCS 1976  $u'$   $v'$  diagram of the chromaticities used in Experiment 1, shown separately for the three types of grating. The bars were generated initially using measurements from a Minolta Color Analyser but the chromaticities shown were measured subsequently using a Photoresearch PR-670 telespectroradiometer; hence the slight variation in chromaticity from the linear pattern expected.

The gratings were horizontal, circular in outline, and from a distance of 0.5m subtended 20 deg of visual angle. A central fixation saltire cross (3mm) was present throughout the entire trial. This was added to provide a stable point for the participant to fixate. Ensuring that the participant was fixating the centre of the grating pattern provided a similar field of view for all patterns for all participants and reduced eye movements that otherwise would have disturbed the autorefractor recording (Wolffsohn, Hunt & Gilmartin, 2002).

### 2.2.3 Procedure

Accommodation was measured using the WAM-5500 autorefractor (manufactured by Grand Seiko Co., Ltd®, Fukuyama, Japan). The autorefractor recorded the accommodative response and the pupil diameter at 2Hz. Recordings were taken from the left eye, but the stimuli were viewed binocularly. The 21 gratings were presented in random order, each for 10 seconds, separated by three seconds during which a grey screen of similar space-averaged luminance was presented. The participant was asked to fixate the central fixation cross throughout the trial. The gratings were presented twice in separate trials separated by a short break.

#### 2.2.4 Data analysis

At the outset of the recording, a measurement of accommodation at 8m was obtained. The spherical value was used to adjust the accommodative response so as to take into account individual refractive error. The mean and standard deviation of participants' accommodative response during the 10 second stimulus presentations were obtained. The mean pupil diameter during stimulus presentation was also analysed.


### 2.3 Results

Data from three participants were excluded because the autorefractor was unable to obtain a reading for all of the stimuli. Poor recordings were due to excessive blinking and/or a small pupil diameter. The accommodative response was expressed as a lag of accommodation by calculating the difference between the accommodative response and the distance of the target in dioptres. The degrees of freedom were corrected for violations in the assumption of sphericity using the Greenhouse Geisser adjustment.

#### 2.3.1 Accommodative lag

Overall, there was no significant correlation between the separation in component chromaticities of the gratings and the mean accommodative lag (Pearson's  $r(19)=-.05, p=.830$ ), or the standard deviation in the accommodative lag ( $r(19)=-.25, p=.273$ ) (Figure 2).

A repeated-measures analysis of covariance with type of grating as factor and colour separation as covariate showed no significant effect of colour separation on the accommodative lag ( $F(1, 153)=0.04, p=.846$ ) and no effect of the grating type (red-blue, blue-green and green-red) ( $F(2, 306)=0.03, p=.968$ ), or an interaction ( $F(2, 306)=1.46, p=.234$ ).




**Figure 2.** On the left, the accommodative lag for each pattern is shown as a function of the colour difference, separately for the three types of pattern (red-blue, blue-green and green-red). The larger numerical value for lag indicates a greater accommodative error. Error bars represent one standard error.

A separate analysis of covariance on the effect of the grating on the standard deviation in the lag, showed that there was no effect of the colour separation ( $F(1,153)=0.90, p=.344$ ), no effect of the colour pair ( $F(1.2,183.4)=3.23, p=.066$ ), or an interaction ( $F(2,306)=1.37, p=.255$ ).

### 2.3.2 Pupil response

The pupil diameter was not significantly related to the type of pattern or the separation in chromaticity of its bars ( $r(19)=-.10, p=.672$ ) (Figure 3).



**Figure 3.** The mean pupil diameter for each pattern is shown as a function of the colour difference, separately for the three types of pattern (red-blue, blue-green and green-red). The error bars represent one standard error.

#### 2.4 Interim discussion

There was no effect of the pattern on accommodative lag, despite a clear effect of the colour difference on discomfort (Haigh et al., 2012b; Wilkins et al., 2008). One possible explanation is that the central fixation cross produced an appropriate luminance contrast for accommodation, which negated any effect of the colour difference on accommodation.

In Experiment 2 we increased the luminance contrast of the pattern. We used the same procedure as Experiment 1, but the patterns were red-black, green-black or blue-black and contained the same central fixation cross. Different wavelengths require different dioptric power due to longitudinal chromatic aberration (LCA). A red object requires more dioptric power compared to a blue object. There is a neurological blur-reducing mechanism that controls the accommodative system (Phillips & Stark, 1977), which indicates that the accommodative response to a red object should be much stronger than the accommodative response to a blue object.

If the results from Experiment 1 were due to the fixation cross, then there should be no difference in the lag between the three patterns. If the fixation cross did not affect accommodation, then the red-black pattern should produce the smallest lag, and the blue-black pattern should produce the largest lag due to LCA.

**Deleted:** 2.3.3 Effect of fatigue on the accommodative lag and pupil diameter¶  
As the recording progressed, there was no significant change in the accommodative lag ( $r(19)=-.06, p=.803$ ) and no change in the standard deviation in the lag ( $r(19)=.03, p=.888$ ). Nevertheless the diameter of the pupil decreased during the recording ( $r(19)=-.94, p<.001$ ) from an average of 4.5mm for the first stimulus to 3.7mm for the last. ¶

**Deleted:** Haigh, Tang, Coutts & Wilkins, submitted

### 3.1 Experiment 2 Effect of LCA on accommodation

#### 3.2 Method

##### 3.2.1 Participants and Stimuli

One male and six females from the University of Essex took part, two of whom participated in Experiment 1. The gratings comprised only red pixels, green pixels or blue pixels, with a space-averaged luminance of 23cd.m<sup>-2</sup> and dimensions similar to those used in Experiment 1.

#### 3.3 Results

A repeated-measures analysis of variance showed that there was a main effect of the colour of the grating (red-black, green-black or blue-black) on the accommodative lag ( $F(2,6)=5.20, p=.049$ ). The blue-black gratings produced the greatest lag compared to both the green-black gratings and the red-black gratings, and the red-black gratings produced a smaller lag than the green-black gratings, but this was not significant when tested using a Bonferroni *post-hoc* test (Table 1). There was also a main effect of the colour of the grating on the standard deviation in the lag ( $F(2,6)=9.21, p=.015$ ). The red-black grating produced the largest standard deviation in the lag compared to the green- and blue-black gratings, but this was not significant when tested using a Bonferroni *post-hoc* test.

**Table 1.** Mean and standard deviation of the accommodative lag from the red-black, green-black and blue-black gratings

Grating	Mean (dioptries)	Standard deviation
Red-black	0.70	0.06
Green-black	0.72	0.05
Blue-black	0.76	0.05

#### 3.4 Interim discussion

There was an effect of LCA on the accommodative lag and the standard deviation in the lag despite the presence of the fixation cross. The effect size of the colour-black gratings (Experiment 2) was much larger ( $\eta_p^2=.634$  mean lag;  $\eta_p^2=.754$  standard deviation of lag) than the effect size of the gratings that varied in their colour separation (Experiment 1) ( $\eta_p^2<.001$  mean lag;  $\eta_p^2=.021$  standard deviation of lag). This suggests that the procedure and instrumentation used in Experiment 1 were sensitive enough to detect changes in accommodative lag if there were changes to be found.

Evidently the findings in Experiment 1 were not due to the fixation cross and the chromatic separation of the bars of the grating did not affect accommodative lag.

If the effects of colour on accommodative lag are attributable to LCA then one might anticipate an effect of LCA on the accommodative response to the large chromaticity separations. The larger the chromaticity separation, the larger the difference in dioptric power needed to de-blur the two bars of the grating. If the accommodative response switches from accommodating to one bar and then accommodating to the other, the large chromaticity separations should have produced a greater variance in the lag. There were, however, no significant effects of chromaticity separation on the variance in the lag detected in Experiment 1 or in Experiment 2.

Atchison *et al.* (2004) compared the variance in the accommodative response to achromatic images and 'multichromatic' images (chromatic Maltise cross on a coloured background). There was no significant difference between the variance in the accommodative response to the multichromatic images and the variance in the response to the achromatic images. The authors suggest that the accommodative system 'favours' accommodating to shorter wavelengths (larger accommodative lags) more than longer wavelengths (smaller accommodative lags).

In Experiment 1, we also found no effect of LCA on the variance in the accommodative lag. This supports the findings of Atchison *et al.* (2004) who showed that the accommodative response does not switch between accommodating to one colour and then another.

In Experiment 1, there appeared to be no differences in accommodative lag between stimuli. This suggests that the aversion to the large colour differences (Haigh *et al.*, 2012b; Wilkins *et al.*, 2008) was not due to a larger accommodative lag or greater variance in the lag.

In the next experiment, the gratings differed with respect to movement. Haigh *et al.* (2012a) reported that aversion from moving gratings was greater than aversion from a static grating. The gratings used here were similar to the gratings used by Haigh *et al.* (2012a); therefore, we can be confident that the moving gratings were more aversive than the static grating. Once again there was no relationship between discomfort and accommodative response.

#### 4.1 Experiment 3 Accommodation to moving gratings

Grating patterns that evoke photoparoxysmal responses in the electroencephalograph in patients with photosensitive epilepsy are generally uncomfortable to view (Wilkins *et al.*, 1984). Gratings

**Deleted:** and nothing to explain the aversion to large chromaticity separations (Wilkins *et al.*, 2008; Haigh, Tang, Coutts and Wilkins, submitted).

**Deleted:** . This absence of any relationship between discomfort and accommodative response was also found in the next set of experiments in which the accommodative response to a different set of uncomfortable grating patterns was measured

that drift at a constant velocity are not epileptogenic, but gratings that drift at the same velocity but reverse direction (creating a vibrating motion) are highly epileptogenic (Binnie, Findlay & Wilkins, 1986). Haigh *et al.* (2012a) reported differences in the contrast thresholds at which drifting, vibrating and static gratings became aversive. The drifting and vibrating gratings were more uncomfortable to view than static patterns. In particular, the drifting gratings (that are not epileptogenic) were perceived to be more uncomfortable than the (highly epileptogenic) vibrating gratings.

One possible explanation for the discrepancy between the gratings that are epileptogenic and those that are aversive is that the accommodative system is unable to accommodate to images that drift at a constant velocity. If this is the case, then we would expect to see a larger accommodative lag and/or more variance in the lag in response to the drifting pattern compared to the static pattern.

The results from Experiment 1 show that there was no effect of the uncomfortable patterns on the accommodative response. If the accommodative response is not responsible for the discomfort from grating patterns, no differences in the lag or the variance in the lag in response to the drifting, vibrating and static patterns would be anticipated.

The accommodative response to the drifting, vibrating and static patterns was measured. The parameters of movements were such as to increase discomfort.

## 4.2 Methods

### 4.2.1 Participants

Four males and 27 females from the University of Essex took part. Twenty nine of these participants also took part in Experiment 1. All participants fulfilled the same criteria as Experiment 1. Participants were asked to complete a questionnaire about their general health, visual history and any headaches they experienced. None of the participants had a history of seizures.

### 4.2.2 Stimuli

Horizontal achromatic gratings were created in MATLAB® using the PsychToolbox extension (Brainard, 1997; Pelli, 1997; Kleiner *et al.*, 2007) with a square-wave luminance profile and a spatial frequency of 2cpd. One of the gratings was made to drift in a vertical direction orthogonal to the grating, at a constant contour velocity of 10 cycles.s<sup>-1</sup>. The second of the gratings had a similar drifting motion, but the movement abruptly reversed direction after one half spatial cycle to create a

vibrating motion. The movement was symmetrical above and below fixation to prevent optokinetic nystagmus; i.e. the movement was downward in the upper visual field and upward in the lower.

The third grating remained static. The gratings were vignettted using a Gaussian window to reduce edge effects. At half contrast the grating subtended 5.6 deg of visual angle. A central fixation cross was superimposed on the gratings (3mm).

Deleted: filter

A uniform grey field ( $54 \text{ cd.m}^{-2}$ ) with a central fixation cross appeared between each grating presentation.

#### 4.2.3 Procedure

Each grating was presented for 10s, followed by 3.5s of grey screen. The central fixation cross (+) was replaced by a black square (side 3mm) 0.5s before the grating was displayed. The gratings were presented in the following order: drifting (D), vibrating (V), stationary(S), S, V, D, S, D, V, V, D, S, V, S, D, S, V, D. Otherwise, the procedure was the same as in Experiment 1.

#### 4.2.4 Data Analysis

The data were analysed as in Experiment 1.

### 4.3 Results

The data from four participants were rejected due the autorefractor being unable to obtain a reading for all of the stimuli, because of excessive blinking or small pupil diameter. As each grating pattern was presented six times, the repetition of the presentations was used as an independent variable in the following analyses of variance.

#### 4.3.1 Accommodative response

A repeated-measures analysis of variance was used to test for differences in the mean (Table 2) and standard deviation of the accommodative response to the three types of grating: drifting, stationary and vibrating. There were no significant differences between the mean ( $F(1.5,36.7)=2.10, p=.133$ ) or the standard deviation ( $F(2,50)=0.47, p=.625$ ) of the accommodative lag to the gratings.

**Table 2.** Mean and standard deviation of the accommodative lag to the drifting, vibrating and static grating patterns.

Grating type	Mean (dioptries)	Standard deviation
Drifting	0.50	0.12
Vibrating	0.54	0.11
Static	0.50	0.12

### 4.3.2 Pupil response

There was no effect of the stimulus type on pupil diameter ( $F(2,50)=0.23, p=.793$ ).

### 4.4 Interim Discussion

There was no effect of the motion of the grating on the accommodative lag or the variance in the lag, indicating that the discomfort from the moving patterns was not due to a larger accommodative lag or more variance in the lag.

Both Experiments 1 and 3 failed to find any association between accommodative lag and the gratings that evoke discomfort. It is therefore difficult to attribute the discomfort to accommodative mechanisms; at least this is the case at the viewing distance used. In the next experiment, we looked at the differences in accommodation to static achromatic patterns and manipulated the perceived viewing distance and thus the accommodative demand. The patterns varied in their spatial frequency and contrast.

## 5.1 Experiment 4 Accommodation to different spatial frequency gratings

Near focal distances place a greater demand on the accommodative system and produce larger lags of accommodation (Morgan, 1968) and more variance in the lag (Jaschinski, 1997). Tosha *et al.* (2009) found that participants with high visual discomfort showed significant signs of fatigue when the distance of the target was 4D compared to 2D. Chase *et al.* (2009) also found larger lags in those with high visual discomfort when viewing targets at 4-5D. If visual discomfort is associated with problems accommodating to the target then increasing the accommodative demand should emphasise any differences in the size of the lag between the patterns that evoke discomfort and those that do not.

**Deleted: ¶**  
**4.3.3 Effect of fatigue on accommodative lag and pupil diameter¶**  
 There was a significant effect of fatigue on mean pupil diameter ( $r(16)=-.81, p<.001$ ). The pupil diameter decreased during the experimental session from an average of 4.0mm to 3.7mm. There was no effect of presentation order on the mean accommodative response ( $r(16)=.20, p=.423$ ) or on the standard deviation ( $r(16)=-.31, p=.216$ )

In this experiment, participants were shown four grating patterns that varied in their spatial frequency and contrast (see Figure 4). The high contrast, mid-range spatial frequency patterns are within the epileptogenic range and are the most uncomfortable to view (Wilkins *et al.*, 1984), so if accommodation is responsible for or related to aversion, we would expect a larger accommodative lag to these patterns. Any effect of the uncomfortable pattern on accommodation should be more apparent when the accommodative demand is greater.

## 5.2 Method

### 5.2.1 Participants

Seventeen females and 20 males from the Leibniz Research Centre for Working Environment and Human Factors (IfADo), Dortmund, Germany participated. Two participants who habitually wore spectacles did so during the study.


### 5.2.2 Stimuli

A horizontal achromatic grating was created and presented using PowerPoint on an 8 inch CTF846-A LED monitor (CarTFT.com, Reutlingen, Germany) powered by a Dell Precision M4500 laptop. A 5D Badal lens was used monocularly to retain the spatial frequency of the grating at various viewing distances. We wished to present monocular accommodative stimuli of constant angular size and identical spatial frequency as a function of viewing distance, i. e. accommodative load. This can be achieved with the Badal optometer principle (Badal, 1876): the eye (i. e. the corneal apex) was placed at the back focal point of a 5 dioptre (Badal) lens, so that the accommodative load (in dioptres) was proportional to the distance of the target from the front focal point of the lens (Atchison *et al.*, 1985): if the target was 20 cm behind the lens, it appeared at optical infinity.

Four achromatic grating patterns were used. All had a square-wave luminance profile and the stripes had a horizontal orientation. All gratings were circular in outline, subtended 8.5deg visual angle, and were surrounded by a grey field of 60 cd.m<sup>-2</sup> space-averaged luminance. A grey fixation dot was superimposed on the grating subtending 0.3deg of visual angle.

The first grating pattern with a spatial frequency of 2cpd was displayed at maximum contrast (>95%) and had a space-averaged luminance of 160 cd.m<sup>-2</sup>. As a comparison for this aversive pattern, we used a version with the same spatial frequency, but with reduced contrast (60%) and reduced space-averaged luminance (85 cd.m<sup>-2</sup>). Given the contribution of contrast and luminance to aversion, this pattern should have been less aversive. The third and fourth patterns had a contrast of >95%

contrast and a [space-averaged](#) luminance of  $160 \text{ cd.m}^{-2}$ , but one had a spatial frequency of 0.25cpd and the other had a spatial frequency of 3.5cpd (Figure 4). The first and fourth patterns had a spatial frequency close to 3 cpd and the contrast and luminance were high and should therefore have been aversive (Wilkins *et al.*, 1984). The second pattern had a lower contrast and luminance and should therefore have been less aversive. The third pattern had a spatial frequency outside the range that induces distortions and discomfort (Wilkins, 1995). The four patterns were presented in the order given above.



**Figure 4.** Schematics of the four grating patterns used in Experiment 4.

### 5.2.3 Procedure

Participants viewed each pattern monocularly, first with one eye and then the other whilst the accommodative response of the viewing eye was measured using an SRW-5000 Autorefractor (Shin-Nippon Commerce Inc: Tokyo, Japan). The patterns were viewed with the left or the right eye first in a counterbalanced order.

For each distance, the participants were asked to “fixieren” (focus on) the central grey fixation [dot](#), whilst single recordings of accommodation were taken (mean 14.5 recordings) over a period of ten seconds. When participants were viewing the gratings monocularly, the other eye was occluded by black cloth which hung from the forehead rest on the autorefractor. The Badal lens was placed 200mm from the viewing eye of the participant. In terms of the accommodative response required, the 200mm distance from the target stimulus to the lens equated to infinity (0D), the 160mm distance to 1m (1D), the 120mm distance to 0.5m (2D), the 80mm distance to 0.33m (3D) and the 40mm distance to 0.25m (4D).


Deleted: cross

### 5.3 Results

The monocular recordings from the left and the right eye were analysed separately, so each participant contributed two recordings to each condition. The accommodative response was converted into accommodative lag by subtracting the accommodative response from the accommodative distance. The lag to the target at different distances (1D – 4D) for each stimulus, and for each eye was subjected to a regression analysis. The average slope for each participant was then calculated over the almost linear range of 1D-4D. Tonic accommodation, which is the resting focal distance, is between 1-1.5D (Leibowitz & Owens, 1978). Analysing the slope of the lag-versus-demand curve allows for greater sensitivity when measuring the effect of a stimulus over various distances.

A large lag and/or a steep lag-versus-demand curve indicates a weak response. This can occur because no vergence or proximal stimuli for accommodation were present. To enhance the sensitivity of the measure, slopes that had a gradient of greater than 0.6 were rejected from the analysis.

In addition, two participants had a negative lag in response to the 1D target consistent with myopia. Following these criteria (>0.6 lag-versus-demand slope, and evidence of myopia), six participants' data were excluded from the analysis. Figure 5 shows the average lag-versus-demand curves for the participants who were included and excluded from the analysis. Note, that including the four participants with slopes that were greater than 0.6 did not affect the findings but did add noise to the results.



Accommodative demand (dioptres)  
Equivalent viewing distance (m)

**Deleted:** (closer than tonic  
**Deleted:** ). In this way, a

**Deleted:** means

**Deleted:** , which


**Deleted:** since

**Deleted:** Response curves

**Deleted:** This was done to ensure that participants were accommodating to the target as instructed. Two

**Deleted:** ,

**Deleted:** <sp>





**Figure 5.** The mean accommodative lag as a function of accommodative demand (equivalent viewing distance) in observers who have been included (left) or excluded (right) from the analysis. The larger the lag, the larger the error in the response. The dotted line shows a slope of 1 (weak accommodative response/large accommodative lag).

The mean and the standard deviation of the accommodative lag was analysed as well as the slope of the lag-versus-demand curve for the different distances of the stimulus.

### 5.3.1 Mean accommodative lag

To ensure that there was an effect of the distance of the stimulus on the accommodative lag, a repeated-measures analysis of variance was conducted. The responses were averaged across stimuli for each distance. There was a significant effect of the accommodative demand ( $F(1.4,42.5)=134.81, p<.001$ ), showing that the near distances created a greater accommodative lag (Figure 6).



**Figure 6.** Accommodative lag as a function of accommodative demand. Error bars showing one standard error.

A paired samples t-test showed that there was no significant difference in the slope of the lag-versus-demand curve between the high and the low contrast gratings ( $t(60)=1.83, p=.072$ ).

A repeated-measures analysis of variance of the slope of the lag-versus-demand curve to the 3.5, 2 and 0.25 cpd gratings showed that there was no effect of the spatial frequency ( $F(1.7,103.2)=2.91, p=.067$ ) on accommodation.

The Accommodative Error Index (AEI) assesses the deviation between the slope of the accommodative response and the slope of the perfect accommodative response (Chauhan & Charman, 1995). The larger the AEI value, the greater the error in the response. The AEI was calculated for the slope from each participant in response to each of the four gratings. There was no significant difference between the high- and low-contrast gratings ( $t(60)=1.08, p=.284$ ), and no effect of the spatial frequency of the grating ( $F(1,60.2)=.97, p=.381$ ).

### 5.3.2 Variance in accommodative response

The standard deviation of the accommodative response was analysed similarly for all stimuli. There was an overall effect of the viewing distance on the variance in the accommodative response ( $F(2.6,160.1)=56.89, p<.001$ ), with the accommodation at nearer distances having a greater standard deviation (Figure 7).





Figure 7. The standard deviation in the accommodative lag as a function of accommodative demand. Error bars representing one standard error.

To analyse the effect of the gratings on the standard deviation of the accommodative lag, analyses of covariance were used, with the accommodative demand as the covariate. When comparing the effect of high and low contrast gratings on the standard deviation in the lag, there was a main effect of the covariate ( $F(1,250)=21.47, p<.001$ ), but there was no effect of the grating ( $F(1,250)=.28, p=.597$ ), and no interaction ( $F(1,250)=.01, p=.918$ ). Similarly, there was no effect of the spatial



- Deleted: Once again,
- Deleted: slope
- Deleted: at the different target distances was calculated and
- Deleted: to test for differences in the variance in accommodation to
- Deleted: different stimuli. A repeated-measures analysis of variance showed that

frequency of the grating ( $F(2,486)=.17, p=.840$ ), and no interaction with the covariate ( $F(2,486)=.12, p=.891$ ).

**Deleted:** pattern

**Deleted:**  $116)=0.11, p=.898$ ). A paired samples t-test showed that there was no difference in the slope of the variance in the accommodative response between the high (2cpd) and low contrast gratings ( $t(60)=-0.31, p=.760$ ).

#### 5.4. Interim Discussion

Once again, there was no effect of the patterns that evoke discomfort on accommodative lag or on the variance in the lag. This was even the case in the 4D condition. Previous studies found that the differences in accommodative lag between the high and low visual discomfort groups were largest when the distance of the target was 4 - 5D. Therefore, if there was an effect of the uncomfortable patterns to be found then this should have been evident in the 4D condition.

It is possible that a longer viewing duration was needed to be able to detect an effect of the grating on the accommodative response. Tosha *et al.* (2009) found effects of fatigue between individuals with high and low visual discomfort after 60 seconds of viewing the stimulus. However, when using the Pattern Glare Test (which measures the number of illusions reported in grating patterns that vary in their spatial frequency) each pattern is only presented to the participant for 3 seconds (Wilkins & Evans, 2001), and yet individuals report discomfort and illusions. The 10 second presentation duration should have therefore been sufficient to detect any effect of uncomfortable stimuli on the accommodative lag.

There was an effect of accommodative demand on the lag and on the standard deviation of the lag. This was expected and is in accordance with previous findings (Morgan, 1968; Jaschinski, 1997). We can therefore be confident that the procedure and the autorefractor used were suitable for detecting differences in accommodation to different targets at a variety of different accommodative distances.

#### 6. General Discussion

For the isoluminant chromatic grating patterns, the moving achromatic patterns, and the patterns that varied in their spatial frequency and contrast, there are clear differences in the discomfort and distortions experienced when viewing the patterns (Haigh *et al.*, 2012a; Haigh *et al.*, 2012b; Wilkins *et al.*, 2008; Wilkins *et al.*, 1984). Nevertheless there was no effect of the pattern parameters on the accommodative lag or on its standard deviation. As participants showed an effect of longitudinal chromatic aberration on the accommodative lag, we cannot conclude that the procedure or the equipment were insufficiently sensitive to measure changes in accommodation, had such changes been associated with discomfort. It is possible that the effect of uncomfortable patterns on

**Deleted:** Haigh, Tang, Coutts &

**Deleted:** , submitted

accommodation is much smaller than the effect of chromatic aberration, but it would be questionable whether such small changes in accommodation could cause the discomfort from the patterns.

The isoluminant patterns that had the greater chromaticity separation would have given rise to greater longitudinal chromatic aberration. The failure to allow for the aberration by adjustment of accommodation might have given rise to discomfort reported by [Haigh et al., \(2012b\)](#) and [Wilkins et al. \(2008\)](#). Such a mechanism would not explain the discomfort from the achromatic patterns, and the absence of any association with accommodative lag or variability in accommodative lag.

Deleted: Haigh

Deleted: submitted

Several studies have found an effect of pattern glare/visual discomfort on accommodation (Allen *et al.*, 2008; Chase *et al.*, 2009; Tosha *et al.*, 2009). Individuals who experience discomfort tend to have a greater lag of accommodation. This appears to be at odds with the findings discussed above. If there is no relationship between accommodation and the patterns that evoke discomfort, then there should be no relationship across individuals between overall visual discomfort and accommodation. It is possible that the larger lag is a mechanism to cope with the discomfort, rather than a cause of the discomfort.

Experiment 4 was conducted monocularly, which could have reduced the discomfort from the patterns. However, Experiment 1 and Experiment 3 were conducted binocularly and there was still no effect of the grating on accommodation.

There is evidence to suggest that the uncomfortable patterns produce a heightened cortical response (Huang *et al.*, 2003; Coutts, Cooper, Elwell & Wilkins, 2012). When the discomfort is reduced by the use of tinted ophthalmic lenses, the cortical response is 'normalised'; that is, the migraineurs then show a similar cortical response to controls (Huang *et al.*, 2011; Coutts *et al.*, 2012). The experiments reported here suggest that the heightened cortical response to the uncomfortable patterns is not dependent on poor accommodation.

There is no indication that accommodation is responsible for the discomfort induced by grating patterns. Any individual differences in accommodative lag are unlikely to drive the discomfort.

## 7. Acknowledgments

We thank those in the Individual Visual Performance group at the Leibniz Research Centre for Working Environment and Human Factors for their help and support. Some of this work was supported by the Wellcome Trust (grant # 80274 to AJW).

## 8. References

- Adjamian, P., Holliday, I. E., Barnes, G. R., Hillebrand, A., Hadjipapas, A., & Singh, K. D. (2004). Induced visual illusions and gamma oscillations in human primary visual cortex. *Eur J Neurosci*, *20*(2), 587-592.
- Allen, P., Hussain, A., Usherwood, C., & Wilkins, A. (2010). Pattern-related visual stress, chromaticity and accommodation. *Invest Ophthalmol Vis Sci*, *51*, 6843-6849.
- Atchison, D. A., Bradley, A., Thibos, L. N., & Smith, G. (1995). Useful Variations of the Badal Optometer. *Optometry & Vision Science*, *72*(4), 279-284.
- Atchison, D. A., Strang, N. C., & Stark, L. R. (2004). Dynamic Accommodation Responses to Stationary Colored Targets. *Optom Vis Sci*, *81*(9), 699-711.
- Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433-436.
- Badal, A. (1876). Nouvel optomètre: Dormant, à la fois et dans une seule opération, la mesure de la réfraction oculaire et celle de l'acuité visuelle. *Ann. ocul.*, *75*(Jan.-Feb.), 1-13.
- Campbell, F. W., & Robson, J. G. (1958). Moving visual images produced by regular stationary patterns. *Nature*, *181*, 362.
- Chase, C., Tosha, C., Borsting, E., & Ridder, W. H. I. (2009). Visual Discomfort and Objective Measures of Static Accommodation. *Optom Vis Sci*, *86*(7), 883-889.
- Chronicle, E. P., & Wilkins, A. J. (1996). Gratings that induce perceptual distortions mask superimposed targets. *Perception*, *25*, 661-668.
- Coutts, L. V., Cooper, C. E., Elwell, C. E., & Wilkins, A. J. (2012). Time course of the haemodynamic response to visual stimulation in migraine, measured using near-infrared spectroscopy. *Cephalalgia*, *32*(8), 621-629.

- Georgeson, M. A. (1976). Psychophysical hallucinations of orientation and spatial frequency. *Perception*, 5(1), 99-111.
- Georgeson, M. A. (1980). The perceived spatial frequency, contrast, and orientation of illusory gratings. *Perception*, 9(6), 695-712.
- Gregory, R. L. (1977). Vision with isoluminant colour contrast: 1. A projection technique and observations. *Perception*, 6(1), 113-119.
- Haigh, S. M., Karanovic, O., Wilkinson, F., & Wilkins, A. J. (2012a). Cortical hyperexcitability in migraine and aversion to patterns. *Cephalgia*, 32(3), 236-240.
- Haigh, S. M., Tang, P., Barningham, L., Coutts, L., Allen, P. M., & Wilkins, A. J. (2012b). Colour separation and aversion. *i-Perception*, 3(4) 232.
- Harle, D. E., Shepherd, A. J., & Evans, B. J. W. (2006). Visual stimuli are common triggers of migraine and are associated with pattern glare. *Headache*, 46, 1431-1440.
- Helmholtz, H. (1856). *Handbuch der Physiologischen Optik*. Leipzig: Leopold Voss.
- Huang, J., Cooper, T., Satana, B., Kaufman, D., & Cao, Y. (2003). Visual distortion provoked by a stimulus in migraine associated with hyperneuronal activity. *Headache*, 43, 664-671.
- Huang, J., Zong, X., Wilkins, A., Jenkins, B., Bozoki, A., & Cao, Y. (2011). fMRI evidence that precision ocular tints reduce cortical hyperactivation in migraine. *Cephalgia*, 31(8), 925-936.
- Jaschinski, W. (1997). Fixation disparity and accommodation as a function of viewing distance and prism load. *Ophthalmic Physiol Opt*, 17(4), 324-339.
- Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3? *Perception*, 36(ECVP Abstract Supplement).
- Liebowitz, H. W. & Owens, D. A. (1978). New evidence for the intermediate position of relaxed accommodation. *Doc Ophthalmol*, 46(1), 133-147.
- Marcus, D. A., & Soso, M. J. (1989). Migraine and Stripe-Induced Visual Discomfort. *Arch Neurol*, 46(10), 1129-1132.

- Millodot, M. (1968). Influence of accommodation on the viewing of an optical illusion. *Q J Exp Psychol*, 20(4), 329-335.
- Morgan, M. W. (1968). Accommodation and vergence. *Am J Optom Arch Am Acad Optom*, 45(7), 417-454.
- Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spat Vis*, 10, 437-442.
- Phillips, S., & Stark, L. (1977). Blur: A sufficient accommodative stimulus. *Documenta Ophthalmologica*, 43(1), 65-89.
- Purkinje, J. (Ed.). (1823). *Beobachtungen und Versuche zur Physiologie der Sinne. Beitrage zur Kenntniss des Sehens in subjectiver Hinsicht* (Erstes Bandchen ed.). Prague: Calve'schen Buchhandlung.
- Purkinje, J. (Ed.). (1825). *Beobachtungen und Versuche zur Physiologie der Sinne. Neue Beitrage zur Kenntniss des Sehens in subjectiver Hinsicht* (Zwites Bandchen ed.). Berlin: Reiner.
- Simmers, A. J., Gray, L. S., & Wilkins, A. J. (2001). The influence of tinted lenses upon ocular accommodation. *Vision Res*, 41, 1229-1238.
- Siniatchkin, M., Moeller, F., Shepherd, A., Siebner, H., & Stephani, U. (2007). Altered cortical visual processing in individuals with a spreading photoparoxysmal EEG response. *Eur J Neurosci*, 26(2), 529-536.
- Switkes, E., Bradley, A., & Schor, C. (1990). Readily visible changes in color contrast are insufficient to stimulate accommodation. *Vision Res*, 30(9), 1367-1376.
- Tosha, C., Borsting, E., Ridder, W. H., & Chase, C. (2009). Accommodation response and visual discomfort. *Ophthalmic Physiol Opt*, 16(6), 625-633.
- Tucker, J., & Charman, W. N. (1987). Effect of target content at higher spatial frequencies on the accuracy of the accommodation response. *Ophthalmic Physiol Opt*, 7(2), 137-142.
- Wade, N. J. (1983). *Brewster and Wheatstone on Vision*. London: Academic Press.
- Wilkins, A. J. (1986). What is visual discomfort? *Trends Neurosci*, 9(8), 343-346.

Wilkins, A. J., & Evans, B. J. W. (2001). *Pattern Glare Test Instructions*. London: i.O.O. Sales Ltd.

Wilkins, A. J., & Nimmo-Smith, M. I. (1987). The clarity and comfort of printed text. *Ergonomics*, 30(12), 1705-1720.

Wilkins, A. J., Nimmo-Smith, M. I., Tait, A., McManus, C., Della Sala, S., Tilley, A., *et al.* (1984). A neurological basis for visual discomfort. *Brain*, 107, 989-1017.

Wolfe, J. M., & Owens, D. A. (1981). Is accommodation colourblind? Focusing chromatic contours. *Perception*, 10, 53-62.

Wolffsohn, J. S., Hunt, O. A., & Gilmartin, B. (2002). Continuous measurement of accommodation in human factor applications. *Ophthalmic Physiol Opt*, 22(5), 380-384.