

BMJ Open Prevalence, determinants and knowledge about herbal medicine and non-hospital utilisation in southwest Nigeria: a cross-sectional study

Olujimi Aina,¹ Lata Gautam ,¹ Padam Simkhada,² Sarah Hall³

To cite: Aina O, Gautam L, Simkhada P, *et al*. Prevalence, determinants and knowledge about herbal medicine and non-hospital utilisation in southwest Nigeria: a cross-sectional study. *BMJ Open* 2020;**10**:e040769. doi:10.1136/bmjopen-2020-040769

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-040769>).

Received 27 May 2020
Revised 04 August 2020
Accepted 07 August 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Life Sciences, Anglia Ruskin University—Cambridge Campus, Cambridge, UK

²School of Human and Health Sciences, University of Huddersfield, Huddersfield, UK

³Leicester School of Pharmacy, Health and Life Sciences, De Montfort University, Leicester, UK

Correspondence to

Dr Lata Gautam;
Lata.Gautam@anglia.ac.uk

ABSTRACT

Objectives To examine the prevalence, determinants, safety perceptions, effectiveness and knowledge of herbal medicines (HMs) and reasons for non-hospital utilisation.

Design Cross-sectional study.

Setting Ekiti state, southwest Nigeria.

Participants A representative sample (n=1600) of adults (18 years or above) currently living in Ekiti state, southwest Nigeria for at least 2 years, at the time of study.

Results The majority of the respondents (85% n=1265) have used HMs in the last 2 years. Across economic classes use, middle income (88.3%) was the highest (p<0.001), suggesting poverty is not a major factor, even with income inequality. Their use was the most common among respondents with a primary level of education (91.4%, p=0.001); and 100% use (p=0.009) of respondents practising African traditional religion; farmers and those 70 years or above. Our study also reveals more men (p<0.001) used HMs (89.9%) than women (78.6%) and effectiveness was a major reason for use (39.6%) followed by affordability (31.9%). Although the majority of the respondents (90%) knew the difference between certified and uncertified HMs, uncertified ones were the most commonly used (37.3%) in the population.

Conclusion Although there is a cultural history of HM use within the study population, the choice of use was based on their effectiveness. Therefore, a scientifically valid analysis of this claim within the study population may help achieve a cheaper and affordable healthcare alternative which will be safe. This is important, considering that uncertified HMs were chosen over certified ones, even though a large majority of respondents were aware of differences and likely consequences. This study highlights the need for further investment by the government, individuals and corporate stakeholders in HM research and improvement of conventional healthcare system. This is in addition to public health awareness on the danger of use of uncertified herbal products.

BACKGROUND

The provision of healthcare falls mainly into the orthodox and non-orthodox systems. The non-conventional healthcare system includes complementary medicine of which herbal medicine (HM) is an important component.¹ HMs are defined as materials or preparations

Strengths and limitations of this study

- Qualitative and quantitative data used in this research catered for the peculiarity of the study population, hence increasing inclusivity.
- Most towns were visited during the day when some farmers may have been absent leading to a low number of participants from this group.
- This study found that the timing of questionnaire administration is an important consideration to make.
- There were more incomplete entries in the self-administered questionnaires compared with the interview-administered ones.

obtained from one or more plants—raw or processed parts²—containing substances with therapeutic characteristics and other benefits to human health. A large percentage of the world's population depends on HMs in both developing (80%)^{3 4} and developed countries.^{5 6}

The use of HM is pivotal to the practice of the African traditional medicine (ATM), and it was the major medical system available to millions of people in urban and rural communities of Africa, until the advent of Europeans.^{7 8} It is a non-conventional system of disease management that employs various processes of consultation with herbalists, priests, media and diverse traditional deities together with herbal use.⁹ The practice of ATM also extended to culturally homogeneous ethnic groups in Yoruba land called the “Ekitis”¹⁰ who form the present Ekiti state. Culturally, the larger Yoruba ethnic group had the Babalawo and Onisekun as traditional medical practitioners and specialists in herbs, respectively, as healthcare providers¹¹ whose existence and practice are still contemporary.

Although, until recently HM in Africa was generally not thoroughly researched and only loosely regulated¹² with the lack of proper documentation of the workings of the ATM



and its practice not helping.¹³ Some of these factors caused the pericolonial conflict between new orthodox systems and the already established ATM, largely due to the belief by the colonialists that the latter was superstitious¹⁴ and therefore declared illegal.¹⁵ The perpetuation of this perception and the lack of proper documentation, research and regulation may still militate against an improved synergy between stakeholders, researchers and collaboration between HM and orthodox medicine. This also includes the general public who are caught between both choices for numerous reasons.

However, recent studies have reported the high use of HMs in Nigeria.¹⁶⁻²⁰ A study conducted in one of the local government areas (Ekiti state) reported 74.3% of the respondents used HM in the treatment of malaria.¹⁶ In another study, in urban Lagos Nigeria, 66.8% (n=388 respondents) used HM for management of various ailments.²⁰ However, these studies are more specific to setting and disease, hence a need for a general population study and non-disease specific study as reported here. Moreover, the knowledge and prevalence of HM use within the larger population in Ekiti state remains unknown, researched or documented; although its use is generally believed to be widespread.

Vendors of various HMs are almost now ubiquitous in Nigeria, trading in traffic gridlocks, highways, bus stops/ stations, festivals and even in some conventional health facilities, but the determinants of its use has not been wholly examined. HMs used are either locally made, refined, imported, certified or uncertified by the government. The National Agency for Food and Drug Administration and Control (NAFDAC) in Nigeria is the government agency responsible for the certification, registration and regulation of HMs. Extemporaneous HMs (herbs or mixture of herbs given on a one-to-one basis) are not included in the class of HMs.²¹ However, some of these products have now been commercialised raising concerns about safety as much as the uncertified HM types, therefore these need to be included in the NAFDAC registration and regulation process.

Patronage of conventional health facilities is an important aspect of healthcare delivery system in most communities. The conventional and traditional medical system provides healthcare services in Ekiti state, coordinated by the State Ministry of Health.²² The increased use of HMs has been attributed to challenges relating to the availability and affordability of conventional medicine. This may be true considering that 40.1% of the total population of Nigeria are poor and live on less than 137430 naira (US\$352) annually.²³

In Ekiti state, of the 16 local government areas, only 2 areas (Irepedun/Ifelodun and Ekiti East) have a higher number of their population using their health facilities optimally.²⁴ However, with the highest poverty head count being 87.7% in Sokoto northwest Nigeria and 4.5% in Lagos southwest Nigeria, Ekiti state has a poverty head count of 28%.²³

Therefore, use of HM is likely a combination of different factors within the study population, hence this

study investigated the knowledge, prevalence and determinant of HM use and the reasons for non-utilisation of conventional health facilities in Ekiti state, Nigeria. The result from this study should help in policy formulation by the government and stakeholders towards improved healthcare access, areas of further research and collaboration. This will be in addition to identification of areas for public health intervention and targeting, which will improve the health outcome of the population.

METHODS

Study design

This was a cross-sectional study conducted in Ekiti state, southwest Nigeria with a population of 2 384 212,²⁵ which include over 127 large and small towns.²⁶ A semi-structured survey was conducted to examine HM use. Participants were 18 years and above and those who lived in Ekiti state at the time of study or have lived in the state for at least 2 years.

The Cochran formula was used for determination of the minimum sample size required in this study. It allows for calculation of the best possible size with preferred precision and confidence level. It is particularly suitable for large study populations and calculates the sample size based on a proportion of people who use HM.²⁷

Hence, the minimum sample size required for this study was 1067 participants assuming 50% of the population use²⁷HM as calculated using equation 1:

$$n = \frac{Z^2 \times pq}{d^2}$$

Where n=required minimum sample size in the study
z=Z-score or SD

p=proportion of people who use HM in the study area
q=1-p

d=the acceptable error level

The calculation of the sample size was based on acceptable error level of 3% (0.03) and a CI of 95% corresponding to a Z-score of 1.96. These are within social science research recommended values for SD and error levels.²⁸ The proportion of people using HM in Ekiti state is unknown, therefore maximum heterogeneity was assumed (ie, a 50/50 split in users and non-users) with p=0.5.

Sampling method

The study participants were selected using a multistage sampling technique. The first stage involved stratifying Ekiti state into the existing 16 local government areas. This followed random sampling to select two towns in each local government (total of 32 towns— second stage). Then residential quarters in each town were selected randomly, making a total of 160 residential quarters (third stage). Systemic random sampling was used in the fourth stage in selecting 10 households from each of the selected residential quarters and one participant from each household made up the final sample population of 1600. Hence, a cross-sectional study was administered to

1600 participants which was above the required minimum sample size as determined earlier and were representatives of the Ekiti state population.²⁵ A flow diagram of the sampling method is shown in online supplemental file 1.

Administration and measurement of research instrument

The questionnaire used in this study was adapted from previous studies on the use of HMs^{18 19 29} and validated in a pilot study. They were interviewer-administered or self-administered depending on the ability or choice of the participant. A total of 10 field enumerators with previous experience were employed from the public and trained to help with data collection. Each self-administered questionnaire was completed in about 10 min and the interviewer-administered in about 20 min. The questionnaire was used to obtain both qualitative and quantitative data with open-ended and closed-ended questions, respectively. Interpretation of the questionnaire in the local dialect was offered when required.

The close-ended questions were used to collect the sociodemographic data of the participants which included their age range, gender, level of education, religion, occupation and annual income. Close-ended questions were also asked to examine knowledge (asked if participants knew what HM was or not, and certification status), their perception of its effectiveness and safety, then the frequency of use among participants to assess prevalence of HM use. The open-ended questions were employed to identify reasons for HM preference and non-orthodox preference. These also included adverse effects experienced and how they were managed.

Data analysis

Manual representational thematic analysis was used to analyse textual data obtained from open-ended questions and presented quantitatively. In some instances where participants gave multiple answers to an open-ended question, each was analysed as individual response. Hence, the total of some responses in a category exceeds the 1265 total number of respondents. Afterwards, inferential and descriptive statistical analysis was carried out using SPSS (V.20.0). Comparison of sociodemographic factors and the use or non-use of HM was carried out using inferential statistics. A Pearson χ^2 test was used to examine the impact of independent variables (eg, age, gender, religion, level of education, occupation, annual income) on various responses (such as use of HM and perception of safety) in the study and $p \leq 0.05$ at 95% CI was considered to be statistically significant. Multicollinearity between the independent variable and the use of HM was also tested. When the variance inflation factor (VIF) and tolerance are greater than 5–10 and lower than 0.1–0.2, respectively, this indicates multicollinearity.

Patient and public involvement

The public have participated in this study by completing the survey. The pilot study was carried out to determine its feasibility on a larger scale. It was observed that some

of the questions on HM use were ambiguous. As a result, the questions were rephrased to focus on the use of HMs within the last 2 years. Research questions were developed from prior knowledge and experience of the researchers. Participants were not asked to assess the burden of the intervention and time required to participate in the research. Multistage sampling method was used for participants' recruitment.

RESULTS

A total of 984 questionnaires were self-administered while 616 questionnaires were interview based. Among these, 1265 respondents formed the final sample size for the study due to exclusion of 335 respondents' entry. A total of 263 self-administered and 72 interview-based questionnaires were excluded due to being incomplete.

Sociodemographic characteristics of respondents

There were more male respondents (56.4%) than female respondents (43.6%) in the study. Majority of the respondents (43.2%) were within the age group of 30–49 years, Christians (69.5%) and self-employed (39.6%). In addition, the majority of the respondents (82%) were within the low-income group as shown in [table 1](#).

Knowledge and use of HMs

Considering their knowledge and use of HMs, 90% of the respondents knew the difference between certified and uncertified HM, while 85% had used them in the last 2 years, 53% using them more than 10 times ([table 2](#)). However, over half of the respondents (57%) believed it is unsafe to take uncertified HM, although 37% continued to take them, while 32% took both certified and uncertified HM.

Preference for HM use and hospital utilisation

The effectiveness was the reason (39.6%) respondents preferred HMs and for those who did not use HMs, risk to health was the major reason (61%). A 45.2% of users blamed poor service delivery for non-hospital utilisation ([table 3](#)).

According to [figure 1](#), across all income classes, effectiveness of HM was the most attributed reason for their use. Likewise, poor service delivery was the most stated reason for non-utilisation of conventional health facilities ([figure 2](#)).

Perception of safety and effectiveness

To explore perception of safety and effectiveness of HMs, our survey results showed that the majority of the respondents (83.6%) perceived that HMs are effective, while more than half (57.3%) believed they are unsafe ([table 4](#)).

Most of the respondents (52.7%) who have used HM in the last 2 years did not experience any adverse effects while some experienced abdominal discomfort (14%), nausea and vomiting (18.7%), headache (7.6%), stooling (5.8%) and generalised body weakness (1.3%). These adverse

Table 1 Sociodemographic background of the respondents

	Frequency	%
Age (years)		
18–29	322	25.5
30–49	547	43.2
50–69	358	28.3
70 and above	38	3.0
Gender		
Male	713	56.4
Female	552	43.6
Level of Education		
No formal education	191	15.1
Primary	245	19.4
Secondary	340	26.9
Tertiary	489	38.7
Religion		
Christianity	879	69.5
Islam	346	27.4
African traditional	40	3.2
Occupation		
Student	84	6.6
Civil servant	491	38.8
Farmer	76	6.0
Self-employed	501	39.6
Others	113	8.9
Annual income (in naira)		
Low (\leq 600 000)	1037	82.0
Middle (600 000–2.4 million)	179	14.2
High (\geq 2.4 million)	49	3.9

Table 2 Knowledge and use of herbal medicines (HMs)


	Frequency	%
Knowledge of certified and uncertified HM		
Have knowledge	1139	90.0
Do not have knowledge	126	10.0
Use of HM in the last 2 years		
Used	1075	85.0
Non-use	190	15.0
Class of HM used		
Uncertified	401	37.3
Both	343	31.9
Certified	331	30.8
Frequency of HM use in last 2 years		
Once–twice	142	13.2
3–10 times	363	33.8
Over 10 times	570	53.0

Table 3 Reasons for herbal medicine (HM) use/non-use and non-hospital utilisation

	Frequency	%
Reasons for HM use		
Effectiveness	538	39.6
Affordability	434	31.9
Availability	292	21.5
Natural product	96	7.1
Reasons for non-HM use		
Risk to health	116	61.1
Personal preference	49	25.8
Poor knowledge of HM composition	25	13.2
Reasons for non-hospital utilisation		
Poor service delivery	534	45.2
High hospital cost	396	33.5
Unorthodox belief	141	11.9
Do visit the hospital	110	9.3

effects were managed differently; the majority (42.5%) took adequate rest while the effects self-limit, visited the hospital (25%), took orthodox medicine (21.6%) or took another HM (10.9%) (see [table 4](#)).

The data have been analysed using Pearson χ^2 test to explore if there is a significant association among different factors studied such as age, sex, level of education, religion, annual income and occupation, and the use of HM. As shown in [table 5](#), results indicate a significant association between, age, sex, religion, annual income and occupation with HM using 95% CI.

**Figure 1** Chart showing reasons for herbal medicine use across economic background of respondents.




Figure 2 Chart showing reasons for non-hospital utilisation across economic background of the respondents.

A test of data for the assumption of collinearity showed multicollinearity was not an issue (age, tolerance=0.86, VIF=1.16; gender, tolerance=0.95, VIF=1.05; level of education, tolerance=0.73, VIF=1.38; religion, tolerance=0.89, VIF=1.29; occupation, tolerance=0.83, VIF=1.21; annual income, tolerance=0.85, VIF=1.17).

DISCUSSION

Various studies have reported the use of HM in health subpopulation³⁰⁻³³ and in different locations and settings.³³⁻³⁵ General population use of HM similar to

this study has been reported in South Africa,³⁶ Benin,³⁷ Uganda³⁸ and Nigeria.^{18 39 40} But the prevalence of HM use in the general population is sparse in sub-Saharan Africa. However, prevalence study on the use of wider traditional complementary medicine reportedly varies from 4.6% in semi-urban Ethiopia⁴¹ to 94% in semi-urban Nigeria.⁴² The variability in prevalence could be a result of factors such as study design and method, socioeconomic variability of sample population, sample size and definition of terms in the study. Likewise the non-utilisation of conventional health system has been attributed to various reasons^{40 43 44}; however, the socioeconomic peculiarity of the study population may also cause differences in findings. Historical and cultural use of HM in Africa presume adequate knowledge of HM already exists; but with the need for its regulation and certification, knowledge testing of the difference between certified and uncertified forms has therefore become more imperative.

In this study the majority of the respondents (90%) knew the difference between government-certified and uncertified HMs. The high level of knowledge about certified and uncertified HMs may have been due to the success of various campaigns and public awareness programmes run by the NAFDAC in 2007 about fake and counterfeit drugs in Nigeria⁴⁵ Participants’ awareness is thus reflected in this study which is an indicator of the effectiveness of the government effort at sensitising the public.

However, we found that the use of HM is still high in the study population with 85% of the respondents having used them in the last 2 years (table 2). This high use has also been described in other African countries such as Ghana, Tanzania, Zambia, Kenya, and about 80% use reported in South Africa.⁴⁶ Similar studies in general HM use conducted in Lagos and the UK reported 66.8% and 64.2% of their respondents had used HM.^{20 47} Social, cultural, socioeconomic and political factors have been reported to influence the use of HM in developed countries,^{47 48} whereas this might not be the case in developing countries such as Nigeria. It has been reported that approximately 80% of Nigerians still consult the traditional healers for various healthcare provisions,⁴⁹ which may explain the high use of HM in this study.

The effectiveness of HMs (83.6%) was a major factor in their use which is in line with previous findings.^{44 50} Affordability was the second most mentioned factor (31.9%) followed by availability (21.5%). These factors influencing HM use in this study have also been reported previously.⁴ In addition, the authors reported other factors such as preference for natural therapies and complementary medicines, dissatisfaction with orthodox pharmaceuticals, high cost and side effects of orthodox medicine, distrust of physician’s abilities and self-medication habit.⁴ Anecdotal information from friends, religious influence and spiritual consciousness have also been known to influence the use of HM.^{51 52} Only 7.1% of the respondents in this study attributed their HM use to its natural and organic properties, the misconception of the natural

Table 4 Perception of herbal medicines’ (HMs) safety and effectiveness and their adverse effect management

	Frequency	%
Perception of effectiveness		
Effective	899	83.6
Ineffective	176	16.4
Perception of safety		
Safe	464	42.7
Unsafe	623	57.3
Observed adverse effect		
None	566	52.7
Abdominal discomfort	150	14.0
Nausea and vomiting	201	18.7
Headache	82	7.6
Stooling	62	5.8
Generalised body weakness	14	1.3
Management of adverse effect		
Rest and becomes self-limiting	219	42.5
Visit the hospital	129	25.0
Take orthodox medicine	111	21.6
Take another HM	56	10.9

**Table 5** Significance of background characteristics on usage of herbal medicines (HMs)

Characteristics	Use of HM	Non-use of HM	Total (100%)	P value
Age (years)				
18–29	274 (85.1%)	48 (14.9%)	322	p=0.005*
30–49	474 (86.7%)	73 (13.3%)	547	$\chi^2=12.996$
50–69	289 (80.7%)	69 (19.3%)	358	df=3
70 and >70	38 (100.0%)	0 (0.0%)	38	
Total	1075 (85.0%)	190 (15.0%)	1265	
Gender				
Male	641 (89.9%)	72 (10.1%)	713	p<0.001*
Female	434 (78.6%)	118 (21.4%)	552	$\chi^2=31.008$
Total	1075 (85.0%)	190 (15.0%)	1265	df=1
Level of Education				
No formal education	169 (88.5%)	22 (11.5%)	191	p=0.001*
Primary	224 (91.4%)	21 (8.6%)	245	$\chi^2=16.670$
Secondary	287 (84.4%)	53 (15.6%)	340	df=3
Tertiary	395 (80.8%)	94 (19.2%)	489	
Total	1075 (85.0%)	190 (15.0%)	1265	
Religion				
Christianity	751 (85.4%)	128 (14.6%)	879	p=0.009*
Islam	284 (82.1%)	62 (17.9%)	346	$\chi^2=9.493$
African traditional	40 (100.0%)	0 (0.0%)	40	df=2
Total	1075 (85.0%)	190 (15.0%)	1265	
Annual Income				
Low income†	890 (85.8%)	147 (14.2%)	1037	p<0.001*
Middle income‡	158 (88.3%)	21 (11.7%)	179	$\chi^2=36.366$
High income§	27 (55.1%)	22 (44.9%)	49	df=2
Total	1075 (85.0%)	190 (15.0%)	1265	
Occupation				
Student	69 (82.1%)	15 (17.9%)	84	p<0.001*
Civil servant	395 (80.4%)	96 (19.6%)	491	$\chi^2=26.701$
Farmer	76 (100.0%)	0 (0.0%)	76	df=4
Business	431 (86.0%)	70 (14.0%)	501	
Others	104 (92.0%)	9 (8.0%)	113	
Total	1075 (85.0%)	190 (15.0%)	1265	

*Significant association.

†(≤600 000 naira).

‡(601 000–2.4 million naira).

§(≥2.4 million naira).

properties being non-toxic and free of adverse effect is a common belief in both developed and developing countries.⁵³ A comparative assessment of herbal and orthodox medicines in Nigeria showed that the HMs were preferred to orthodox medicines in terms of efficacy, affordability, availability, safety and level of advertisement.⁴⁰

The dissatisfaction with orthodox healthcare was also highlighted in this study. Majority of the respondents (45.2%) attributed poor service delivery in the hospital to non-hospital utilisation followed by high hospital

cost (33.5%) (table 3). The poor services experienced included long hospital waiting time, hostile and rude attitude of health workers, poor infrastructure and lack of communication. The nature of illness, availability of money, patient's age, religious or personal belief, level of education and severity of illness among other factors have been reported to influence the choice between orthodox and the traditional health system.

Nonetheless, challenges associated with the orthodox health system in Ekiti state such as affordability of service

cost, quality of service rendered, closeness to home, staff attitude, environment cleanliness, availability of required drugs and services have also informed patient's patronage to the use of HM.⁵⁴ In addition, approximately 70% of orthodox drugs in circulation in Nigeria are either adulterated or fake highlighting the need for thorough analysis of such products and licensing provisions.

Hence, findings from this study suggest lack of improvement in the healthcare service delivery over time, which is also evidenced by other publications highlighting similar issues.⁵³ The effectiveness of HM and hospital poor service delivery are the reasons most highlighted in this study favouring HM use which is across the whole socioeconomic class of the respondents (figures 1 and 2). Thus, regardless of the economic situation of the respondents, a good service delivery in the orthodox health system is highly desirable.

A significant difference in the number of men using HM (89.9%) was observed when compared with women (78.6%) ($p < 0.001$, $\chi^2 = 31.008$ and $df = 3$) (table 5). Our finding is not consistent with published literature from developed countries where women had higher use of HM than men with 12.7% of women compared with 1.1% of men⁵⁵ and 8.3% women compared with 2.9% of men⁵⁶ with a similar trend reported in other studies.^{57 58} A previous Nigerian study however found no significant difference between being women and use of HM.⁵⁹ Although it has been reported that women take less risk than men,⁶⁰ women tend to seek help from the orthodox healthcare when compared with men.^{60–63} Therefore, we recommend authorities to be mindful of targeting men in public health awareness in relation to the safety of HM.

The level of education was another factor that contributed to the use of HM in this study. There was significant relationship between respondent's level of education and use of HM ($p = 0.001$, $\chi^2 = 16.670$, $df = 3$), as respondents with primary level of education had the highest use of HM and least use within respondents with tertiary level of education. HM use was also highest among respondents within the middle annual income class while the least use was among the high annual income group ($p < 0.001$, $\chi^2 = 36.366$, $df = 2$). This finding contradicts a previous study where the author has reported poverty and dissatisfaction with orthodox medicine as the reason for HM use.⁶⁴ In line with the author, our study showed dissatisfaction with orthodox medical care as a reason for non-hospital utilisation (table 3) but poverty was not a factor.⁶⁴ Affordability which was the second most highlighted reason for use of HM (31.9%) and high hospital cost which was the second most highlighted reason for non-hospital utilisation (33.5%) may not equate to poverty. Though 82% of the respondents were within the low-income class, 69% of healthcare financing is still 'out of pocket' by patients in Nigeria.⁶⁵ Health insurance accounts for only 2% of all healthcare financing in Nigeria.⁶⁴

Therefore, financing out-of-pocket healthcare is an extra burden on a population with 82% already within the low-income group. This brings to bare the importance

of the United Nations Sustainable Development Goal 3, which is to strive towards achieving Universal Health Coverage by 2030 by all member states.⁶⁶

Conversely, low level of education and income power has been reported to influence choice of HM use in previous studies.^{18 57 67} The studies associated low income with the use of HM, which is different from what was found in this study. Highest use of HM was among the middle-income group (table 5). However, studies in developed countries have reported a non-significant relationship between income and HM use but a significant relationship between higher level of education and HM use.^{29 68} These variations may be due to interplay between factors such as respondent's socioeconomic characteristics and peculiarities of the study environment. Hence, the findings in this study reflect such interplay where the middle-income class used more HM and a significant relationship between level of education and HM use.

Additionally, there was a significant association between the age of the respondents and use of HM in this study ($p = 0.005$, $\chi^2 = 12.996$, $df = 3$). Respondents 70 years and above have all (100%) used HM in the last 2 years, followed by the age group 30–49 years of which 86.7% of them have also used HM within the period. A publication among the general population on HM use reported significant relationship between age and use of HM,⁵⁷ while another reported no significant difference.¹⁸ This study showed that the use of HM increases with the age of the respondents, although age group 50–69 years did not follow this pattern. The occupation of the respondents was also significantly associated with HM use ($p = 0.001$, $\chi^2 = 75.504$ $df = 8$). This can be linked to the annual income of the respondents which had significant association in this study. The type of occupation largely determines the accruable income.

There was 100% use of HM in the last 2 years among respondents who practised African traditional religion, 85.4% use among Christians and 82.1% use among Muslims. There was a significant relationship between religious affiliation of the respondents and HM use ($p = 0.009$, $\chi^2 = 9.493$, $df = 2$) (table 5). The use of HM as an integral part of African traditional religion is documented and African traditional religionists being the custodian of the African traditional medicine.⁶⁹ Although some studies reported no significant association between religion and HM use,^{36 39 50} other studies have reported otherwise.^{70 71} Furthermore, research has shown that religious beliefs and spiritual practices of patients have powerful influence in making decision about treatment choice, coping with chronic diseases and end-of-life care decisions.^{72 73} Therefore, this study highlights the influence religion has on the use of HM; it is apparently a correlation between their religious belief and their choice of healthcare. There is no multicollinearity between the sociodemographic characteristics and the use of HM, therefore it reduces standard errors in related independent variables such as level of education, education and annual income.



This study employed qualitative and quantitative data which catered for the peculiarity of the study population, hence increasing inclusivity. Also, the study has helped to reveal the prevalence and determinant of HM use and non-hospital utilisation within the study population. These are new findings within the Ekiti state which will help in public health planning. There were few limitations of this study. First, there were more incomplete entries in the self-administered questionnaires compared with the interview-administered ones, which reduced the number of eventual participants in the study. Second, most of the towns were visited during the day, at which farmers would have been away from the residential areas. This led to a low number of participants from this group compared with other occupations in this study. However, the eventual sample size was still representative of the study population. Also, there were possibilities for recall bias due to respondents' need to remember answers related to the previous 2 years, but the methodology used in this study is acceptable and robust for contemporary research in public health.

Further research may be needed to assess the use of HM in subhealth and the general population to increase present knowledge on HM use and ways to integrate traditional and orthodox healthcare systems within the study population.

CONCLUSIONS

Findings from this study showed majority of the study population have used HMs in the last 2 years regardless of the economic class. Majority of the respondents in the study know the difference between certified and uncertified HMs, but uncertified HMs were most commonly used. This study highlights the need for further investment in public health enlightenment and healthcare system. This is important considering the findings from this study on reasons for non-hospital utilisation corroborating the health infrastructural/service deficit which has been reported to influence the use of HMs in developing countries. However, the choices are limited; a choice between an acclaimed effective HM and an orthodox health system offering unsatisfactory service. While affordability was also an important reason people used HMs in this study, high hospital cost was also a reason people did not patronise orthodox medical service. It is a choice between an affordable HM and an expensive orthodox medical service especially in the absence of adequate health insurance coverage. This study has shown that there is a significant association between annual incomes, education, gender, age, religion and occupation with the use of HMs. Although there is a cultural history of HM use within the study population, socioeconomic and sociodemographic factors were determinants of its use in light of a prevalent income inequality and absence of adequate health insurance coverage.

Twitter Lata Gautam @LataGautam10 and Sarah Hall @SarahHa81702781

Contributors OA designed and implemented the work; carried out data analysis and interpretation; drafted the manuscript; and is accountable for the accuracy and integrity of the work. LG designed and supervised the work; carried out interpretation; drafted and revised the manuscript for final approval; and is accountable for the accuracy and integrity of the work. PS advised on public health aspect of the research and carried out interpretation and revision of the manuscript. SH supervised the work; carried out interpretation; drafted and revised the manuscript for final approval; and is accountable for the accuracy and integrity of the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethics approval was obtained from the Faculty of Science and Technology, Anglia Ruskin University and Ekiti State Ministry of Health in Ekiti state. Participants were provided with the participant information sheet and the content explained in English or local dialect. Informed consent was obtained from the participants who volunteered to be part of the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. All data relevant to the study are included in the article.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Lata Gautam <http://orcid.org/0000-0002-0409-4724>

REFERENCES

- 1 Frass M, Strassl RP, Friehs H, *et al*. Use and acceptance of complementary and alternative medicine among the general population and medical personnel: a systematic review. *Ochsner J* 2012;12:45–56.
- 2 World Health Organization (WHO). Traditional complementary and integrative medicine. Available: <https://www.who.int/traditional-complementary-integrative-medicine/about/en/> [Accessed 03 Nov 2019].
- 3 Bodeker C, Bodeker G, Ong CK, *et al*. *WHO global atlas of traditional, complementary and alternative medicine*. Geneva, Switzerland: WHO, 2005.
- 4 Bandaranayake WM. Quality Control, Screening, Toxicity, and Regulation of Herbal Drugs. In: *Modern Phytomedicine. Turning medicinal plants into drugs*. Weinheim: Wiley-VCH GmbH & Co. KGaA, 2006: 25–57.
- 5 Nissen N. Practitioners of Western herbal medicine and their practice in the UK: beginning to sketch the profession. *Complement Ther Clin Pract* 2010;16:181–6.
- 6 Calapai G. European legislation on herbal medicines: a look into the future. *Drug Saf* 2008;31:428–31.
- 7 Romero-Daza N. Traditional medicine in Africa. *Ann Am Acad Pol Soc Sci* 2002;583:173–6.
- 8 Abdullahi AA. Trends and challenges of traditional medicine in Africa. *Afr J Tradit Complement Altern Med* 2011;8:115–23.
- 9 MacGaffey W. African traditional religion. Oxford bibliographies. Available: <http://www.oxfordbibliographies.com/view/document/obo-9780199846733/obo-9780199846733-0064.xml#firstMatch> [Accessed 25 Jul 2020].
- 10 Oguntomisin GO. Political change and adaptation in Yorubaland in the nineteenth century. *Can J Afr Stud* 1981;15:223–37.
- 11 Olayiwola DO. The Aladura: It's Strategy for Mission and Conversion in Yoruba land, Nigeria. *Orita: Ibadan J Rel Stud* 1987;19:40–56.
- 12 Mills E, Cooper C, Seely D, *et al*. African herbal medicines in the treatment of HIV: Hypoxis and Sutherlandia. An overview of evidence and pharmacology. *Nutr J* 2005;4:19.
- 13 Van Wyk BE, Van Oudtshoorn B, Gericke N. *Medicinal plants of South Africa*. 336. Pretoria: Briza Publications, 1999. ISBN: 978-1-875093-37-3.
- 14 Onwuanibe RC. The philosophy of African medical practice. *Issue: A Journal of Opinion* 1979;9:25–8.

- 15 Helwig D. Traditional African medicine. Gale encyclopedia of alternative medicine. Available: <http://www.encyclopedia.com/doc/1G2-3435100785.html> [Accessed 24 Jul 2020].
- 16 Olorunniyi OF, Morenikeji OA. The extent of use of herbal medicine in malaria management in Ido/Osi local government area of Ekiti state, Nigeria. *J Med Plants Res* 2013;7:3171–8.
- 17 Awodele O, Amagon KI, Usman SO, Kennedy IA, Sikiru OU, *et al.* Safety of herbal medicines use: case study of ikorodu residents in Lagos, Nigeria. *Curr Drug Saf* 2014;9:138–44.
- 18 Okoronkwo I, Onyia-Pat J-L, Okpala P, *et al.* Patterns of complementary and alternative medicine use, perceived benefits, and adverse effects among adult users in Enugu urban, Southeast Nigeria. *Evid Based Complement Alternat Med* 2014;2014:1–6.
- 19 Duru CB, Uwakwe KA, Chinomso NC, *et al.* Socio-Demographic determinants of herbal medicine use in pregnancy among Nigerian women attending clinics in a tertiary hospital in Imo state, South-East, Nigeria. *Am J Med* 2016;4:1–10.
- 20 Oreagba IA, Oshikoya KA, Amachree M. Herbal medicine use among urban residents in Lagos, Nigeria. *BMC Complement Altern Med* 2011;11:117.
- 21 NAFDAC guidelines herbal and cosmetics. Available: <https://www.nafdac.gov.ng/resources/guidelines/> [Accessed 20 Jul 2020].
- 22 Ekiti state government strategic health development plan 2010–2015, 2011. Available: <https://drive.google.com/file/d/0B1DAMtM1BcbMVXpvRlhxaUdYOFU/view> [Accessed 20 Jul 2020].
- 23 National Bureau of Statistics (NBS). Poverty and inequality in Nigeria: Executive summary. available at, 2019. Available: <https://nigerianstat.gov.ng/download/1092> [Accessed 20 Jul 2020].
- 24 Odeyemi CA. Spatial distribution of poverty indicators in Ekiti state. *Dev Ctry Stud* 2014;4:119.
- 25 National Population Commission. Census report, 2006. Available: <https://nigeriacrvs.gov.ng> [Accessed 03 Nov 2019].
- 26 Ekiti state government EKSG, 2016. Available: <https://ekitistate.gov.ng/about-ekiti/> [Accessed 03 Nov 2019].
- 27 Cochran WG. *Sampling techniques*. 3rd Ed. New York: Wiley, 2007.
- 28 Bryman A, Cramer D. *Quantitative data analysis for social scientists*. London, New York: Routledge, 1990.
- 29 Kummert CM, Spector ML, Dawson DV, *et al.* Patterns of complementary and alternative medicine (CAM) use among dental patients. *J Public Health Dent* 2015;75:109–17.
- 30 Olusanya BO, Inem VA, Abosede OA. Infants delivered in maternity homes run by traditional birth attendants in urban Nigeria: a community-based study. *Health Care Women Int* 2011;32:474–91.
- 31 Sarmiento I, Zuluaga G, Andersson N. Traditional medicine used in childbirth and for childhood diarrhoea in Nigeria's cross river state: interviews with traditional practitioners and a statewide cross-sectional study. *BMJ Open* 2016;6:e010417.
- 32 Girma E, Tesfaye M. Patterns of treatment seeking behavior for mental illnesses in Southwest Ethiopia: a hospital based study. *BMC Psychiatry* 2011;11:138.
- 33 Awodele O, Agbaje EO, Abiola OO, *et al.* Doctors' attitudes towards the use of herbal medicine in Lagos, Nigeria. *J Herb Med* 2012;2:16–22.
- 34 Mbutho NP, Gqaleni N, Korporaal CM. Traditional complementary and alternative medicine: knowledge, attitudes and practices of health care workers in HIV and AIDS clinics in Durban hospitals. *Afr J Tradit Complement Altern Med* 2012;9:64–72.
- 35 van Staden AM, Joubert GBA. Interest in and willingness to use complementary, alternative and traditional medicine among academic and administrative university staff in Bloemfontein, South Africa. *Afr J Tradit Complement Altern Med* 2014;11:61–6.
- 36 Hughes GD, Aboyade OM, Clark BL, *et al.* The prevalence of traditional herbal medicine use among hypertensives living in South African communities. *BMC Complement Altern Med* 2013;13:38.
- 37 Allabi AC, Busia K, Ekanmian V, *et al.* The use of medicinal plants in self-care in the Agonlin region of Benin. *J Ethnopharmacol* 2011;133:234–43.
- 38 Auerbach BJ, Reynolds SJ, Lamorde M, *et al.* Traditional herbal medicine use associated with liver fibrosis in rural Rakai, Uganda. *PLoS One* 2012;7:e41737.
- 39 Opara ER, Osayi KK. Factors affecting the utilization of herbal medicine as a Livelihood alternative among residents of IMO state: the role of social work professionals. *IOSR-JHSS* 2016;21:66–78.
- 40 Osemene KP, Elujoba AA, Ilori MO. A comparative assessment of herbal and orthodox medicines in Nigeria. *Res J. Med Sci* 2011;5:280–5.
- 41 Flatie T, Gedif T, Asres K, *et al.* Ethnomedical survey of Berta ethnic group Assosa zone, Benishangul-Gumuz regional state, mid-west Ethiopia. *J Ethnobiol Ethnomed* 2009;5:14.
- 42 Sarki ZM, Danjuma MH. Socio-Demographic factors and utilization of traditional medicine in Kazaure town, Jigawa state, Nigeria. *Int J Emerg Know* 2015;3:9–20.
- 43 Aikins I, Ahmed M, Adzimah ED. Assessing the role of quality service delivery in client choice for healthcare: a case study of Bechem government hospital and green Hill Hospital. *Eur J Purch Supply Manag* 2014;2:1–23.
- 44 Adesiji GB, Komolafe SE. Factors influencing the use of traditional healing among farmers in Kwara state, Nigeria. *Albanian J Agric Sc* 2013;12:275–81.
- 45 Ofuani AB, Kuye OL, Ogundele OJK. An assessment of the efficiency of government regulatory agencies in Nigeria. Case of the National agency for food and drugs administration and control. *Acad J Econ Stud* 2015;1:9–26.
- 46 Ozioma EOJ, Nwamaka OA. Herbal medicines in African traditional medicine, 2019. Available: <https://www.intechopen.com/books/herbal-medicine/herbal-medicines-in-african-traditional-medicine> [Accessed 20 Jul 2020].
- 47 Ong P, Bodeker G. Use of complementary and alternative medicine services in England. *Am J Public Health* 2002;92:1653–6.
- 48 Eskinazi D. Factors that will Shape the Future of Alternative Medicine: An Overview. In: *What will influence the future of alternative medicine? A world perspective*. Singapore: World Scientific Publishers, 2001: 1–22.
- 49 Adefolaju T. Traditional and orthodox medical systems in Nigeria: the imperative of a synthesis. *AJHR* 2014;2:118–24.
- 50 Mafimisebi TE, Oguntade AE. Health Infrastructure Inequality and Rural-Urban Utilization of Orthodox and Traditional Medicines in Farming Households: A Case Study of Ekiti State, Nigeria. In: Smigorski K, ed. *Health Management - Different Approaches and Solutions*. In Tech, 2011: 197–214.
- 51 Astin JA. Why patients use alternative medicine: results of a national study. *JAMA* 1998;279:1548–53.
- 52 Zeil H. Complementary alternative medicine boon or boondoggle. *Skeptic* 1999;7:86–9.
- 53 Wertheimer Al, Wang PG. *Counterfeit medicines. policy, economics and counter measures*. First Edition. ILM Publications USA, 2012: 1–3.
- 54 Omotoso D. Health seeking behaviour among the rural dwellers in Ekiti state, Nigeria. *Afr. Res. Rev.* 2010;4:125–38.
- 55 Harrison RA, Holt D, Pattison DJ, *et al.* Who and how many people are taking herbal supplements? A survey of 21,923 adults. *Int J Vitam Nutr Res* 2004;74:183–6.
- 56 Raji MA, Kuo Y-F, Snih SA, *et al.* Ethnic differences in herb and vitamin/mineral use in the elderly. *Ann Pharmacother* 2005;39:1019–23.
- 57 Gardiner P, Graham R, Legedza ATR, *et al.* Factors associated with herbal therapy use by adults in the United States. *Altern Ther Health Med* 2007;13:22–9.
- 58 Bakhotmah BA, Alzahrani HA. Self-Reported use of complementary and alternative medicine (CAM) products in topical treatment of diabetic foot disorders by diabetic patients in Jeddah, Western Saudi Arabia. *BMC Res Notes* 2010;3:254.
- 59 Onyeka TC, Ezike HA, Nwoke OM, *et al.* Herbal medicine: a survey of use in Nigerian presurgical patients booked for ambulatory anaesthesia. *BMC Complement Altern Med* 2012;12:130.
- 60 Harris CR, Jenkins M. Gender differences in risk assessment. why do women take fewer risks than men? *Judgm Decis Mak* 2006;1:48–63.
- 61 Green CA, Pope CR. Gender, psychosocial factors and the use of medical services: a longitudinal analysis. *Soc Sci Med* 1999;48:1363–72.
- 62 Bertakis KD, Azari R, Helms LJ, *et al.* Gender differences in the utilization of health care services. *J Fam Pract* 2000;49:147.
- 63 Chang H-J, Lai Y-L, Chang C-M, *et al.* Gender and age differences among youth, in utilization of mental health services in the year preceding suicide in Taiwan. *Community Ment Health J* 2012;48:771–80.
- 64 Oshikoya KA, Senbanjo IO, Njokanma OF, *et al.* Use of complementary and alternative medicines for children with chronic health conditions in Lagos, Nigeria. *BMC Complement Altern Med* 2008;8:66.
- 65 Uzochukwu BSC, Ughasoro MD, Etiaba E, *et al.* Health care financing in Nigeria: implications for achieving universal health coverage. *Niger J Clin Pract* 2015;18:437–44.
- 66 Sustainable Development Goal (SDG). Sustainable development knowledge platform. Available: <https://sustainabledevelopment.un.org/sdg3> [Accessed 20 Jul 2020].
- 67 Barnes PM, Bloom B, Nahin RL. Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report* 2008;10:1–23.



- 68 Du Y, Wolf I-K, Zhuang W, *et al.* Use of herbal medicinal products among children and adolescents in Germany. *BMC Complement Altern Med* 2014;14:218.
- 69 White P. The concept of diseases and health care in African traditional religion in Ghana. *HTS Teologiese Studies / Theological Studies* 2015;71:1–7.
- 70 Hulela K, Thobega M. Botswana Faith Leaders' Attitudes towards the Use of Indigenous Medicinal Herbs. *J Med Plants Res* 2013;7:1550–6.
- 71 Nicdao EG, Ai AL. Religion and the use of complementary and alternative medicine (cam) among cardiac patients. *J Relig Health* 2014;53:864–77.
- 72 Puchalski CM. Spirituality and health: the art of compassionate medicine. *Hospital Physician* 2001:30–6.
- 73 McCormick TR, Hopp F, Nelson-Becker H, *et al.* Ethical and spiritual concerns near the end of life. *J Relig Spiritual Aging* 2012;24:301–13.