


# BMJ Open Internet-based cognitive-behavioural therapy for tinnitus: secondary analysis to examine predictors of outcomes

Hansapani Rodrigo,<sup>1</sup> Eldré W Beukes,<sup>2,3</sup> Gerhard Andersson,<sup>4,5,6</sup> Vinaya Manchaiah <sup>2,7</sup>

**To cite:** Rodrigo H, Beukes EW, Andersson G, *et al.* Internet-based cognitive-behavioural therapy for tinnitus: secondary analysis to examine predictors of outcomes. *BMJ Open* 2021;**11**:e049384. doi:10.1136/bmjopen-2021-049384

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online. (<http://dx.doi.org/10.1136/bmjopen-2021-049384>).

Received 26 January 2021  
Accepted 09 August 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Vinaya Manchaiah;  
[vinaya.manchaiah@lamar.edu](mailto:vinaya.manchaiah@lamar.edu)

## ABSTRACT

**Objectives** The current study examined predictors of outcomes of internet-based cognitive-behavioural therapy (ICBT) for individuals with tinnitus.

**Design** Secondary analysis of intervention studies.

**Setting** Internet-based guided tinnitus intervention provided in the UK.

**Participants** 228 individuals who underwent ICBT.

**Interventions** ICBT.

**Primary and secondary outcome measures** The key predictor variables included demographic, tinnitus, hearing-related and treatment-related variables as well as clinical factors (eg, anxiety, depression, insomnia), which can have an impact on the treatment outcome. A 13-point reduction in Tinnitus Functional Index (TFI) scores has been defined as a successful outcome.

**Results** Of the 228 subjects who were included in the study, 65% had a successful ICBT outcome. As per the univariate analysis, participants with a master's degree or above had the highest odds of having a larger reduction in tinnitus severity (OR 3.47; 95% CI 1.32 to 12.51), compared with the participants who had education only up to high school or less. Additionally, the baseline tinnitus severity was found to be a significant variable (OR 2.65; 95% CI 1.50 to 4.67) contributing to a successful outcome with the intervention. Both linear and logistic regression models have identified education level and baseline tinnitus severity to be significant predictor variables contributing to a reduction in tinnitus severity post-ICBT. As per the linear regression model, participants who had received disability allowance had shown a 25.3-point lower TFI reduction compared with those who did not experience a decrease in their workload due to tinnitus after adjusting for baseline tinnitus severity and their education level.

**Conclusions** Predictors of intervention outcome can be used as a means of triaging patients to the most suited form of treatment to achieve optimal outcomes and to make healthcare savings. Future studies should consider including a heterogeneous group of participants as well as other predictor variables not included in the current study.

**ClinicalTrials.gov Registration:** NCT02370810 (completed); NCT02665975 (completed)

## INTRODUCTION

Tinnitus is the perception of sounds in the absence of external stimulation and is often heard as a ringing or buzzing meaningless sound(s). It is a very common condition with

## Strengths and limitations of this study

- The current study, to our knowledge, is the first study to use combined data from multiple studies to examine the predictors of internet-based cognitive-behavioural therapy (ICBT) outcome for tinnitus.
- The study included a homogeneous group of tinnitus patients due to the strict inclusion/exclusion criteria and may not have included all the possible variables (eg, health literacy, acceptability and motivation of users, satisfaction from the intervention, intervention engagement) that may have played a role in ICBT outcome.
- The sample size remained relatively small when compared with the number of predictive factors included which may have hampered the study results.
- The multivariable analyses may have some limitations in terms of examining complex relationships. Other statistical models including artificial intelligence and machine learning techniques may have more value in examining the non-linear relationship.

at least 15% of the adult population having tinnitus.<sup>1</sup> Tinnitus is highly heterogeneous, both in the way it manifests as well as in the manner those with tinnitus respond to treatment options.<sup>2</sup> The National Study of Hearing in England found that of the general population surveyed (N=48, 313), 10.1% reported any tinnitus, 2.8% reported moderately annoying tinnitus, 1.6% reported severely annoying tinnitus and 0.5% were unable to lead a normal life due to the severity of the tinnitus.<sup>1</sup> Although there are several management strategies described in the literature, most are not evidence based. The main exception is cognitive-behavioural therapy (CBT), as indicated in various systematic reviews of randomised controlled trials.<sup>3-6</sup> Clinical practice guidelines based on research evidence and expert consensus recommend CBT as a management option for individuals with tinnitus and is supported by the American Academy of Otolaryngology-Head and Neck Surgery.<sup>7</sup>

Despite positive and strong evidence for CBT, individuals with tinnitus are rarely offered CBT in their local clinics. For example, a large-scale epidemiological study in the USA showed that medication (which is not recommended for tinnitus in clinical guidelines) was discussed 50% of the time by health professionals, whereas the CBT was discussed as a management option only 0.2% of the time.<sup>8</sup> This is most likely a consequence of the limited number of trained professionals who provide CBT for tinnitus. One solution to overcome this issue is to use Internet-based CBT (ICBT), in which patients are provided with CBT in a self-help format over the internet with minimal guidance from a tinnitus expert.<sup>9</sup> A series of controlled studies in Sweden, Germany the UK and the USA have demonstrated positive effects of ICBT in reducing tinnitus distress as well as reducing its comorbidities such as anxiety, depression and insomnia.<sup>9</sup> In addition to the changes noted in standardised outcome measures, the qualitative analysis of user experiences has highlighted the perceived benefits of this programme.<sup>10</sup> In addition, the improvements noted from ICBT have been maintained for 1-year postintervention.<sup>11</sup> These results suggest that ICBT is a highly promising approach to provide evidence-based tinnitus management.

Although the previous studies on ICBT have shown favourable results, group effects were mainly reported. There is limited understanding of who is likely to benefit (or not) from the ICBT intervention. In other words, only a few studies have examined predictors of ICBT outcomes in tinnitus research. For example, the long-term analysis of the previous UK studies suggested that the best predictors of tinnitus improvements at 1 year were the baseline tinnitus severity, engagement with ICBT programme (ie, more modules read) and higher self-reported satisfaction with the intervention.<sup>11</sup> Studies in other health areas have also examined the predictors of outcome for a range of internet-based health interventions.<sup>12–16</sup> These studies have inconsistently identified various demographic as well as disease-specific variables that could predict the successful and non-successful participants on internet interventions.<sup>16</sup> There remains a clear gap in knowledge in terms of predictors of ICBT outcomes for tinnitus.

Predictors of intervention outcomes may help triage patients to the most suitable tinnitus intervention. If interventions are recommended based on their suitability, it can potentially improve the outcomes resulting in health-care savings. The objective of the current study was to examine the predictors of outcomes of ICBT intervention for individuals with tinnitus based on the secondary analysis of the pooled results from the three-phase clinical trial undertaken in the UK.

## METHOD

### Study design and participants

A large data set was sought to identify predictors of outcome. Trials with similar methodologies were hence sought to merge to form a larger data set. Although a

few previous studies regarding ICBT were conducted in Europe, extensive outcome measures were not used. Following these trials, three trials were conducted in the UK using the same outcome measures. These trials were used due to a lack of other controlled trials available to pool data from. This present study, thus, formed a secondary analysis of data collected from three separate ICBT trials. Study participants from the three separate trials with different designs including the single-group pretest and post-test design,<sup>17</sup> an efficacy RCT design (ClinicalTrials.gov: NCT02370810),<sup>18</sup> and an effectiveness RCT design (ClinicalTrials.gov: NCT02665975)<sup>19</sup> were combined. These studies were conducted during 2016–2018. In the efficacy trial, the experimental group underwent ICBT immediately after allocation whereas the control group underwent the same intervention following an 8-week weekly check-in period. In the effectiveness trial, the experimental group underwent the ICBT intervention whereas the control group underwent treatment as usual, consisting of in-person tinnitus counselling and sound therapy, provided by the audiologists at three hospital settings. The data were collected from only those who underwent the ICBT intervention and were included in this study. The study team was granted access to the deidentified datasets, not containing any personally identifiable information, as part of a data sharing policy. The study results are presented using the Transparent Reporting of a multivariable prediction model for Individual Prognosis or Diagnosis checklist (see online supplemental file 1).

Combining the data from three trials resulted in the inclusion of 228 participants. Of these, 36 were from the pilot trial, 146 were from the efficacy trial and the remaining 46 were from the effectiveness trial.

### Intervention

The intervention included a CBT programme that was specifically developed for individuals with tinnitus.<sup>20</sup> This intervention was originally developed by psychologists in Sweden,<sup>21</sup> but later adapted by audiologists in the UK<sup>22</sup> and the USA.<sup>23</sup> The intervention was administered using a secured ePlatform<sup>24 25</sup> and presented in a self-help format. The intervention was presented over an 8-week period, during which the users were given access to 2–3 modules each week. The CBT programme was divided into 21 modules, of which five were optional. The modules included content such as applied relaxation, thought analysis, cognitive restructuring, imagery and exposure techniques. Each module included text, images and videos to enhance the user experience. In addition, users were required to complete various exercises to engage them in the intervention. Although the intervention was presented in a self-help format, the users had access to minimal guidance from an audiologist (EWB). Generally, this included examining weekly exercises users completed and providing feedback as well as answering any questions they may have in the secured messaging system. An average of 10 min per participant was spent

on providing guidance and support, although some users required more support.

### Outcome measures

The study participants completed an extensive pre-intervention questionnaire that collected data on demographics, tinnitus-related and treatment-related history. In addition, participants also completed various standardised patient-reported outcome measures at baseline (T0), at postintervention (T1) and at the 2 month follow-up (T2). The primary outcome measure included the Tinnitus Functional Index (TFI)<sup>26</sup> to assess tinnitus severity/distress. This is a 25-item questionnaire with scores ranging between 0 and 100. Scores below 25 indicate mild tinnitus with no need for intervention, scores ranging between 25 and 50 indicate a significant problem with possible need for intervention, and scores above 50 indicate severe enough tinnitus possibly requiring a more intensive intervention. The TFI has good psychometric properties with acceptable internal consistency (0.97) and test-retest reliability (0.8).<sup>26</sup>

The secondary outcome measures included the Insomnia Severity Index<sup>27</sup> as a measure of insomnia, the generalised anxiety disorder<sup>28</sup> as a measure of anxiety, the Patient Health Questionnaire<sup>29</sup> as a measure of depression, the Hearing Handicap Inventory for Adults Screening version<sup>30</sup> as a measure of self-reported hearing disability, the Hyperacusis Questionnaire<sup>31</sup> to assess the presence hyperacusis (ie, reduced tolerance of everyday sounds), the Cognitive Failures Questionnaire<sup>32</sup> was administered to assess cognitive functions, and the Satisfaction with Life Scales<sup>33</sup> to assess the global life satisfaction.

### Patient and public involvement

As a secondary analysis, no patients were involved in these studies. The data originate from individuals with tinnitus who had previously received CBT delivered via the internet (ie, ICBT). As the same protocol was followed for all study participants and all received the same intervention, merging this data was possible.

### Variables included in the predictive model

#### Outcome variable

The dependent variable was the pre-and post-intervention change in tinnitus distress based on the TFI score (TFI change). The 13-point change in TFI scores identified as a clinically meaningful (or significant) change by the original authors<sup>26</sup> was used to define a clinically significant intervention outcome.

#### Predictor variables

Predictor variables were selected based on clinical reasoning and findings from previous studies by Beukes *et al*<sup>1</sup> (see online supplemental file 2 for details). Thirty-two variables were selected as potential predictor (independent) variables and included demographic, tinnitus and hearing-related variables, tinnitus treatment related variables. Clinical factors are as follows:

- ▶ **Demographic variables (n=7):** age (dichotomous), gender (dichotomous), education level (ordinal), employment type (categorical), loud noise exposure (dichotomous), diagnosed with a psychological condition (dichotomous), work less due to tinnitus (categorical).
- ▶ **Tinnitus and hearing-related variables (n=15):** baseline tinnitus severity (dichotomous), tinnitus duration (dichotomous), how often tinnitus heard (ordinal), tinnitus location (categorical), tinnitus types (nine different types, dichotomous), multiple tones heard (dichotomous) and hearing loss (categorical).
- ▶ **Treatment-related to tinnitus (n=4):** past treatment sought (dichotomous), sounds can distract from tinnitus (ordinal), hearing aid use (categorical) and medication use (dichotomous).
- ▶ **Clinical factors (n=7):** anxiety (dichotomous), depression (dichotomous), insomnia (dichotomous), hyperacusis (dichotomous), hearing disability (dichotomous), cognitive functions (dichotomous) and life satisfaction (dichotomous).

### Data analysis

The data were analysed using descriptive statistics as well as univariate and multivariable linear regression and logistic regression models. Linear models were used to identify the factors affecting a significant TFI score change while the logistic model was used to evaluate the factors which specifically effect outcomes and was thus selected. There were 98 subjects who had all their predictive variables except their post TFI scores. With the intention of preserving the power of the analysis, we have retained those subjects in the analysis after applying the predictive mean matching (PMM) data imputation.<sup>34</sup> Data imputation with PMM has been identified to be less vulnerable to model misspecification as there is no need to define an explicit model for the distribution of the missing values.<sup>35</sup>

The univariate analysis was performed using  $\chi^2$  or Fisher's exact test to examine the effect of single variables on the ICBT outcome using all the variables. The multivariable regression model was used to identify the effect of the variables on tinnitus reduction post ICBT while adjusting for the baseline tinnitus severity as a variable previously identified to relate to the success of ICBT.<sup>11</sup> Prior to the multivariable analyses, the full data set was divided into training (80%, n=183) and testing (20%, n=45) to make a fair comparison among all the predictive models. The training data set was used to develop the corresponding multivariable regression models while the testing data set was used to evaluate the model predictions. Several competing multivariable models (both linear and logistic) were examined. The best models were selected based on the lowest mean squared error and the lowest Akaike information criterion (AIC).<sup>36</sup> During multivariable analysis, we began with the full model, including all the predictor variables and used backward elimination based on AIC to select the final model. R squared and adj. R squared values have been reported, as they are

**Table 1** Details of clinical variables of the study participants

Characteristic	Mean (SD)
Preintervention tinnitus severity (measured using TFI, scores range 0–11)	57.93 (19.17)
Postintervention tinnitus severity (measured using TFI, scores range 0–11)	34.22 (22.76)
2-month follow-up tinnitus severity (measured using TFI, scores range 0–11)	34.23 (24.19)
Anxiety (measured using GAD-7, scores range 0–21)	7.29 (5.52)
Depression (measured using PHQ-9, scores range 0–27)	7.61 (5.73)
Insomnia (measured using ISI, scores range 0–28)	12.49 (6.67)
Hyperacusis (measured using HQ, scores range 0–40)	18.33 (9.05)
Hearing disability (measured using the HHIA-S, scores range 0–40)	16.18 (11.64)
Cognitive failures (measured using the CFQ, scores range 0–100)	38.54 (15.63)
Life satisfaction (measured using SWLS, scores range 0–40)	20.71 (7.55)

CFQ, Cognitive Failures Questionnaire; GAD-7, generalised anxiety disorder; HHIA-S, Hearing Handicap Inventory for Adults Screening; HQ, Hyperacusis Questionnaire; ISI, Insomnia Severity Index; PHQ-9, Patient Health Questionnaire; SWLS, Satisfaction with Life Scales; TFI, Tinnitus Functional Index.


statistical measures of fit that indicate how much variation of the outcome is explained by the predictor variable(s) in a linear regression model.<sup>37</sup> We also reported the mean squared error as it is a better measure of prediction accuracy. Both crude and model-based ORs were calculated and used to evaluate the effect of the variable. The Hosmer-Lemeshow goodness-of-fit statistic was calculated to assess the calibration of the final model.<sup>38</sup>

The dependent variable TFI change was used as a continuous variable for a linear regression analysis whereas the dichotomous variable (ie, 13-point change yes or no) was used for logistic regression analysis. All statistical analyses were performed with R statistical software (V.3.6.3). All tests were two sided and threshold at 5% level of significance.

## RESULTS

### Participant demographics

The mean age of study participants was 55.14 (SD: 12.92) years, and 57% of the subjects (n=130) were males. The mean tinnitus duration was 17.68 (SD: 19.42) years. Further details on demographic, tinnitus, hearing-related and treatment-related variables are provided in online supplemental file 2 (tables 1–4). **Table 1** presents details on clinical variables. The mean baseline tinnitus severity and tinnitus severity following the ICBT intervention were



**Figure 1** Tinnitus severity (TFI scores) preintervention and postintervention. Boxplot represents the five-number summary (minimum, first quartile, median, third quartile and maximum). The thick dark line represents the median. TFI, Tinnitus Functional Index.

57.93 (SD: 19.17) and 34.22 (SD: 22.76), respectively. **Figure 1** presents the preintervention and postintervention tinnitus severity (TFI) score variation, indicating statistically significant differences between these scores ( $p < 0.001$ ) with the paired t-test. There were 148 participants (65%) with a 13-point or higher reduction after the intervention.

### Univariate analysis to examine the predictors of ICBT outcome

With the exception of education level ( $p = 0.01$ ), none of the demographic variables were associated with post-intervention tinnitus severity change of 13 points or more. Participants with a master's degree or above had the highest odds of having a larger severity change score, with an OR of 3.47 (95% CIs 1.32 to 12.51), compared with the participants who had education only up to high school or less. In terms of tinnitus and hearing-related variables, the baseline tinnitus severity ( $p = 0.001$ ) was significantly associated with treatment success. Participants who had a higher baseline tinnitus severity (ie, TFI scores of greater than or equal to 55.2) had significantly higher odds of treatment success (OR 2.65; 95% CIs 1.50 to 4.67) compared with those who had a baseline severity less than 55.2. The details of the univariate analyses are provided in online supplemental file 3 (tables 1–5).

In terms of the treatment-related variables, sounds can distract ( $p = 0.001$ ) showed a significant association with treatment success. Those who reported being distracted by the sound partially (OR 4.34; 95% CI 1.82 to 10.34) or not at all (OR 3.15; 95% CIs 0.99 to 10.00) were at higher odds of having a successful treatment outcome when compared with those who were fully distracted. However, the odds among the participants who used hearing aids either in one ear or both ears compared with those who did not were not statistically significantly different with a  $p = 0.26$  (see online supplemental table 4). Tinnitus described as voice-like had a 91% lower odds of success

**Table 2** The best multiple linear regression model summary

Predictor variable	Estimate	95% CI	P value
Intercept	-28.94	-41.70 to 16.18	<0.0000
Work less: no	Ref		
Work less: reduced hours	-6.25	-23.90 11.39	0.48
Work less: stopped work	-0.58	-10.52 to 9.36	0.91
Work less: disability allowance	-25.30	-46.35 to 4.24	<b>0.02</b>
Baseline tinnitus severity	0.83	0.65 to 1.00	<b>&lt;0.0001</b>
Education level: high school or less	Ref		
Education level: college	-2.25	-12.61 to 8.11	0.67
Education level: vocational training	0.98	-10.29 to 12.25	0.86
Education level: bachelor's degree	5.14	-4.13 to 14.42	0.28
Education level: master's degree or above	16.81	5.78 to 27.84	<b>0.003</b>

CI, Confidence Interval

with the treatment. None of the clinical factors were significantly associated with the outcome.

#### Multivariable analyses to examine predictors of ICBT outcome

Working less due to tinnitus ( $p=0.046$ ), baseline tinnitus severity ( $p<0.001$ ) and education level ( $p=0.014$ ), showed significant associations with outcome (ie, TFI reduction). Modified models with the remaining variables were not statistically significant. Moreover, several two-way interactions were tested. We did not find any gender interactions with regard to the maskability of sounds ( $p=0.87$ ) and) and hearing aid usage ( $p=0.68$ ) variables. The overall model resulted with an R squared=0.35 and Adj. R squared of 0.20. The final model resulted in a root mean square of 22.81 on the testing data set. All required regression assumptions were satisfied with the selected model. The final regression model (see [table 2](#)) was selected with backward elimination based on AIC.

This model indicated that those who received disability allowance due to having severe tinnitus and being unable

to work had shown a reduction of 25.30 points (95% CIs -46.35 to -4.24) in TFI compared with those who did not have to work less due to tinnitus. Moreover, for every 10 unit increase in the baseline tinnitus severity, there was a 8.3 points (95% CIs 0.65 to 1.00) reduction in their TFI score after adjusting for other variables. Participants who had master's degree or above compared with participants who had a college education showed an expected reduction of 17 points (95% CIs 5.78 to 27.84) in their TFI score.

Multivariable logistic regressions were performed next and indicated that baseline tinnitus severity ( $p<0.001$ ) and education level ( $p=0.001$ ) were identified as significant predictors (see [table 3](#)). This model had an AIC of 212.21. Modified models to the prior model indicated that other variables were not statistically significant (see [table 4](#)).

The multivariable model adjusted OR (see [table 3](#)) for the participants who had master's level or above

**Table 3** The multivariable logistic regression model summary and the model adjusted OR (95% CI) for successful ICBT outcome of 13 points of higher

	Estimate	P value	Model-based adjusted OR (95% CI for OR)
Intercept	-2.32	0.0005	0.10 (0.03 to 0.37)
Baseline tinnitus severity	0.04	<b>&lt;0.001</b>	<b>1.04 (1.02 to 1.06)</b>
Education level: high school or less	Ref		
Education level: college	-0.4	0.41	0.67 (0.26 to 1.74)
Education level: vocational training	0.41	0.47	1.49 (0.50 to 4.48)
Education level: bachelor's degree	0.68	0.14	1.98 (0.79 to 4.98)
Education level: master's degree or above	2.27	<b>0.001</b>	<b>9.65 (2.32 to 40.15)</b>

CI, Confidence Interval

; ICBT, internet-based cognitive-behavioural therapy; OR, Odds Ratio

**Table 4** Predictor variables which were insignificant in multivariable regression models

	Predictor variable	P value	
		Multivariable linear regression model	Multivariable logistic regression model
1	Gender	0.47	0.83
2	Hearing loss	0.89	0.72
3	Tinnitus type: ringing	0.38	0.91
4	Tinnitus type: buzzing	0.43	0.53
5	Tinnitus type: high pitch	0.56	0.48
6	Tinnitus type: low pitch	0.33	0.46
7	Tinnitus type: pulsing	0.99	0.34
8	Tinnitus type: clicking	0.09	0.01
9	Tinnitus type: music	0.37	0.69
10	Tinnitus type: voices	0.34	0.09
11	Tinnitus type: humming	0.96	0.06
12	Anxiety	0.07	0.48
13	Depression	0.76	0.86
14	Insomnia	0.94	0.53
15	Hyperacusis	0.75	0.53
16	Hearing disability	0.84	0.57
17	Cognitive functions	0.71	0.72
18	Life satisfaction	0.75	0.84
19	Multiple tones heard	0.26	0.81
20	Loud noise exposure	0.32	0.76
21	Work less due to tinnitus	Refer <a href="#">table 2</a>	0.46
22	Presence of a psychological condition	0.88	0.72
23	Past treatment sought	0.60	0.83
24	Hearing aid use	0.21	0.20
25	Sounds can distract	0.51	0.11
26	Medication use	0.73	0.87
27	Tinnitus location	0.50	0.27
28	Employment type	0.63	0.90
29	Age	0.88	0.70
30	Tinnitus duration	0.17	0.93
31	How often tinnitus is heard	0.23	0.57

education compared with those who had high school education or less also showed 9.65 higher odds (95% CIs 2.32 to 40.15) of having a successful outcome. Similar to the linear regression model, baseline tinnitus severity had also shown a significant association (OR 1.04; 95% CIs 1.02 to 1.06) with the treatment outcome. The Hosmer-Lemeshow goodness-of-fit test confirmed a better fit in the current model with a  $p$ -value of 0.50 ( $\chi^2=7.36$ ,  $df=8$ ).

Fewer variables were statistically significant in the logistic regression model, which identified influencing predictors of the ICBT success. This was due to the fact that the logistic regression model evaluated predictors of treatment successes (ie, 13-point change), while the

multivariable regression model identified the predictors of a significant TFI reduction.

## DISCUSSION

Accessible and affordable tinnitus interventions are needed to alleviate tinnitus distress as well as comorbid problems with anxiety, depression and insomnia. The current study examined predictors of outcomes for ICBT. In this exploratory study, only a limited number of variables were identified as possibly reducing tinnitus severity scores on the TFI by at least 13 points following ICBT intervention, and the results vary depending on the model

used. Only educational level and baseline tinnitus severity were predictors in both linear and logistic models. The other significant variable in the linear regression models included the demographic variable, work restrictions due to tinnitus when controlling for baseline tinnitus severity and education level. These key findings are discussed below.

In terms of demographic variables, education level was found to be a significant predictor of ICBT success as those with a master's education or higher had higher odds of having a successful outcome compared with those with high school education in both the linear and logistic models. This was expected as having good literacy skills is essential when understanding the intervention materials. The intervention materials used in these studies were written at an average of ninth-grade reading level<sup>23</sup> suggesting that they were not easily accessible for participants with only a high school education. These results highlight the importance of health literacy considerations when developing text-based self-help interventions such as ICBT. Additionally, those who reported work restrictions due to tinnitus were at a lower odds of having a successful outcome. This finding needs further exploration in future studies. Working may, for instance, provide some distraction from tinnitus as supported by reports during the 2020 COVID-19 pandemic that tinnitus was more bothersome for some individuals due to the lack of distractions from commuting and sounds at work.<sup>39</sup> Closely monitoring the effects of tinnitus is important to ensure that tinnitus can be managed so that individuals are still able to work effectively.

When examining the tinnitus and hearing-related variables, baseline tinnitus severity was found to be a significant predictor of ICBT success, as seen in previous studies.<sup>11</sup> Tinnitus perceptions vary greatly, and in this study, those with tinnitus presenting as musical, lower-pitched or clicking were less likely to have a positive outcome of ICBT. This finding certainly needs further exploration as the limited number of participants in each group of tinnitus perception. One of the CBT intervention aims is to help participants to reinterpret their tinnitus to a less threatening sound. It may be that these sounds are not easily likened to everyday sounds than other types of tinnitus (ie, buzzing, high pitch, pulsing, humming) making it difficult to develop adaptation strategies.

Of the four treatment-related variables, only those who reported to use of wearing one hearing aid were found to be at better odds of ICBT success. This finding needs further exploration to identify other characteristics that may be associated with an outcome such as having tinnitus in only one ear. Although the evidence for the use of hearing aids alone for tinnitus management is limited,<sup>40 41</sup> hearing aids may for some reduce the tinnitus percept and aid communication difficulties.<sup>42</sup> Ensuring hearing loss is addressed in addition to the provision of ICBT may lead to more optimal outcomes for those with coexisting hearing loss.

Regarding studying the clinical factors, those with higher levels of depression were found to have a higher reduction in the TFI score. However, the participants with insomnia showed lower odds of success. Interestingly, other clinical factors including anxiety, hyperacusis, hearing disability as well as cognitive functioning were not significant predictors of ICBT in the current study. Further studies and models are required to verify these results.

Studies in other health areas have also examined the predictors of a range of internet-based health interventions.<sup>12–15</sup> Generally, higher baseline symptoms predict increased treatment response, as in anxiety and depression,<sup>43</sup> and higher obsessive-compulsive behaviours when treating the obsessive-compulsive disorder.<sup>44</sup> Variables such as age and gender have been mentioned as significant predictors for some ICBT interventions.<sup>15 43</sup> Most previous ICBT interventions have not identified pretreatment characteristics to predict or moderate outcomes.<sup>16</sup> Most ICBT studies have indicated that ICBT works irrespective of treatment history.<sup>43</sup> Contrarily, previous treatment has shown worse outcomes in some previous studies.<sup>45</sup> However, it may be that some participants may have sought alternative therapies which have no evidence for tinnitus. For this reason, it would be useful to examine specific types of previous treatments in future studies to distinguish between those who had evidence-based interventions before enrolling to ICBT than those who did not.

### Study limitations and future research

The current study was to our knowledge the first study to combine data from multiple studies to examine the predictors of ICBT outcome for tinnitus. However, it has limitations. First, the study may have included a homogeneous group of tinnitus patients due to study inclusion/exclusion criteria and may not have included all the possible variables (eg, health literacy, acceptability and motivation of users, satisfaction from the intervention, intervention engagement) that may have played a role in ICBT outcome. These factors were not investigated for this study. As they have been found to contribute to outcomes,<sup>46</sup> they should be included in future studies. Second, the sample size remained relatively small when compared with the number of predictive factors included. Third, multivariable analyses may have some limitations in terms of examining complex relationships. Moreover, due to the high multicollinearity between the predictor variables, there were several competing models which had led to the same prediction accuracies and root mean square errors. Additionally, these linear models lack in identifying any predictor variables that have a non-linear relationship with the response variables. For these reasons, the study results must be viewed as preliminary. Future studies may benefit from using non-linear statistical models such as generalised additive models, and also artificial intelligence and machine learning models like neural

networks, random forest and support vector machines, as some variables like tinnitus duration and depression had shown lower correlation with the response (with correlations:  $-0.10$  and  $0.29$ , respectively). In addition, including more relevant predictive factors (eg, health literacy, motivation, engagement, adherence) in future studies may help improve predictive accuracy. Currently, we have used AIC value to compare the competing models. For future studies, we are planning to use average Area Under the Receiver Operating Characteristic curve (AUC) and Brier scores to compare models.

#### Author affiliations

<sup>1</sup>School of Mathematical and Statistical Sciences, The University of Texas Rio Grande Valley, Edinburg, Texas, USA

<sup>2</sup>Department of Speech and Hearing Sciences, Lamar University, Beaumont, Texas, USA

<sup>3</sup>Department of Vision and Hearing Sciences, Anglia Ruskin University, Chelmsford, UK

<sup>4</sup>Department of Behavioral Sciences and Learning, Linköping University, Linköping, Sweden

<sup>5</sup>Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden

<sup>6</sup>Department of Clinical Neuroscience, Division of Psychiatry, Karolinska Institute, Stockholm, Sweden

<sup>7</sup>Department of Speech and Hearing, School of Allied Health Sciences, Manipal University, Manipal, Karnataka, India

**Twitter** Eldré W Beukes @Eldre7, Gerhard Andersson @profGergardA and Vinaya Manchaiah @VManchaiah

**Contributors** HR, VM, EWB and GA conceptualised the study. EWB administered the intervention and collected the data. HR performed the data analysis. VM and HR prepared the first draft of the manuscript. All authors read and approved the final manuscript.

**Funding** This work was partially funded by the National Institute on Deafness and Communication Disorders (NIDCD) of the National Institute of Health (NIH) under the award number R21DC017214.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The ethical clearance for these studies was obtained from the Faculty of Science and Technology Research Ethics Panel of Anglia Ruskin University (ARU reference: FST/FREP/14/478 and FST/FREP/14/478) and the East of England–Cambridge South Research Ethics Committee (REC reference: 16/EE/0148) and Health Research Authority (IRAS project ID: 195565).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Vinaya Manchaiah <http://orcid.org/0000-0002-1254-8407>

#### REFERENCES

- Davis A, Refaie AE. The epidemiology of tinnitus. In: Tyler R, ed. *The Handbook of tinnitus*. Singular, 2020: 1–23.
- Cederroth CR, Gallus S, Hall DA, et al. Editorial: towards an understanding of tinnitus heterogeneity. *Front Aging Neurosci* 2019;11:53.
- Hesser H, Weise C, Westin VZ, et al. A systematic review and meta-analysis of randomized controlled trials of cognitive-behavioral therapy for tinnitus distress. *Clin Psychol Rev* 2011;31:545–53.
- Hoare DJ, Kowalkowski VL, Kang S, et al. Systematic review and meta-analyses of randomized controlled trials examining tinnitus management. *Laryngoscope* 2011;121:1555–64.
- Landry EC, Sandoval XCR, Simeone CN, et al. Systematic review and network meta-analysis of cognitive and/or behavioral therapies (CBT) for tinnitus. *Otol Neurotol* 2020;41:153–66.
- Fuller T, Cima R, Langguth B, et al. Cognitive behavioural therapy for tinnitus. *Cochrane Database Syst Rev* 2020;38:CD012614.
- Tunkel DE, Bauer CA, Sun GH, et al. Clinical practice guideline: tinnitus. *Otolaryngol Head Neck Surg* 2014;151:S1–40.
- Bhatt JM, Lin HW, Bhattacharyya N. Prevalence, severity, exposures, and treatment patterns of tinnitus in the United States. *JAMA Otolaryngol Head Neck Surg* 2016;142:959–65.
- Beukes EW, Manchaiah V, Allen PM, et al. Internet-Based interventions for adults with hearing loss, tinnitus, and vestibular disorders: a systematic review and meta-analysis. *Trends Hear* 2019;23:233121651985174.
- Beukes EW, Manchaiah V, Davies ASA, et al. Participants' experiences of an Internet-based cognitive behavioural therapy intervention for tinnitus. *Int J Audiol* 2018c;57:947–54.
- Beukes EW, Allen PM, Baguley DM, et al. Long-Term efficacy of Audiologist-guided Internet-based cognitive behavior therapy for tinnitus. *Am J Audiol* 2018d;27:431–47.
- Blankers M, Koeter MWJ, Schippers GM. Baseline predictors of treatment outcome in Internet-based alcohol interventions: a recursive partitioning analysis alongside a randomized trial. *BMC Public Health* 2013;13:455.
- Chen H, Rodriguez MA, Qian M, et al. Predictors of treatment outcomes and adherence in Internet-based cognitive behavioral therapy for social anxiety in China. *Behav Cogn Psychother* 2020;48:291–303.
- Jonas B, Tensil M-D, Leuschner F, et al. Predictors of treatment response in a web-based intervention for cannabis users. *Internet Interv* 2019;18:100261.
- Spek V, Nyklíček I, Cuijpers P, et al. Predictors of outcome of group and Internet-based cognitive behavior therapy. *J Affect Disord* 2008;105:137–45.
- Andersson G, Titov N, Dear BF, et al. Internet-Delivered psychological treatments: from innovation to implementation. *World Psychiatry* 2019;18:20–8.
- Beukes EW, Allen PM, Manchaiah V, et al. Internet-Based intervention for tinnitus: outcome of a single-group open trial. *J Am Acad Audiol* 2017;28:340–51.
- Beukes EW, Baguley DM, Allen PM, et al. Audiologist-guided Internet-based cognitive behavior therapy for adults with tinnitus in the United Kingdom: a randomized controlled trial. *Ear Hear* 2018;39:423–33.
- Beukes EW, Andersson G, Allen PM, et al. Effectiveness of guided Internet-based cognitive behavioral therapy vs face-to-face clinical care for treatment of tinnitus: a randomized clinical trial. *JAMA Otolaryngol Head Neck Surg* 2018;144:1126–33.
- Beukes E, Andersson G, Manchaiah V. *Cognitive behavioral therapy for tinnitus*. San Diego, USA: Plural Publishing Inc, 2021.
- Andersson G, Strömberg T, Ström L, et al. Randomized controlled trial of Internet-based cognitive behavior therapy for distress associated with tinnitus. *Psychosom Med* 2002;64:810–6.
- Beukes EW, Vlaescu G, Manchaiah V, et al. Development and technical functionality of an Internet-based intervention for tinnitus in the UK. *Internet Interv* 2016;6:6–15.
- Beukes EW, Fagelson M, Aronson EP, et al. Readability following cultural and linguistic adaptations of an Internet-based intervention for tinnitus for use in the United States. *Am J Audiol* 2020;29:97–109.
- Vlaescu G, Alasjö A, Miloff A, et al. Features and functionality of the Iterapi platform for Internet-based psychological treatment. *Internet Interv* 2016;6:107–14.



- 25 Manchaiah V, Vlaescu G, Varadaraj S, *et al.* Features, functionality, and acceptability of Internet-based cognitive behavioral therapy for tinnitus in the United States. *Am J Audiol* 2020;29:476–90.
- 26 Meikle MB, Henry JA, Griest SE, *et al.* The tinnitus functional index: development of a new clinical measure for chronic, intrusive tinnitus. *Ear Hear* 2012;33:153–76.
- 27 Bastien CH, Vallières A, Morin CM. Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Med* 2001;2:297–307.
- 28 Spitzer RL, Kroenke K, Williams JBW, *et al.* A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166:1092–7.
- 29 Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. primary care evaluation of mental disorders. patient health questionnaire. *JAMA* 1999;282:1737–44.
- 30 Newman CW, Weinstein BE, Jacobson GP, *et al.* Test-Retest reliability of the hearing handicap inventory for adults. *Ear Hear* 1991;12:355–7.
- 31 Khalfa S, Dubal S, Veuillet E, *et al.* Psychometric normalization of a hyperacusis questionnaire. *ORL J Otorhinolaryngol Relat Spec* 2002;64:436–42.
- 32 Broadbent DE, Cooper PF, FitzGerald P, *et al.* The cognitive failures questionnaire (CFQ) and its correlates. *Br J Clin Psychol* 1982;21:1–16.
- 33 Diener E, Emmons RA, Larsen RJ, *et al.* The satisfaction with life scale. *J Pers Assess* 1985;49:71–5.
- 34 Lodder P. *To impute or not impute: That's the question. Advising on research methods: Selected topics.* Huizen: Johannes van Kessel Publishing, 2013.
- 35 Little RJA, Rubin DB. *Statistical analysis with missing data.* New York: John Wiley & Sons, 2002.
- 36 Akaike H. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 1974;19:716–23.
- 37 Montgomery DC, Peck EA, Vinning GG. *Introduction to linear regression analysis.* Wiley Series in Probability and Statistics, 2012.
- 38 Agresti A. *Categorical data analysis.* 3rd edn. Hoboken: John Wiley & Sons Inc, 2013.
- 39 Beukes EW, Baguley DM, Jacquemin L, *et al.* Changes in tinnitus experiences during the COVID-19 pandemic. *Front Public Health* 2020;8:592878.
- 40 Hoare DJ, Edmondson-Jones M, Sereda M, *et al.* Amplification with hearing AIDS for patients with tinnitus and co-existing hearing loss. *Cochrane Database Syst Rev* 2014;50:CD010151.
- 41 Shekhawat GS, Searchfield GD, Stinear CM. Role of hearing AIDS in tinnitus intervention: a scoping review. *J Am Acad Audiol* 2013;24:747–62.
- 42 Del Bo L, Ambrosetti U. Hearing AIDS for the treatment of tinnitus. *Prog Brain Res* 2007;166:341–5.
- 43 Stjerneklar S, Hougaard E, Thastum M. Guided Internet-based cognitive behavioral therapy for adolescent anxiety: predictors of treatment response. *Internet Interv* 2019;15:116–25.
- 44 Andersson E, Ljótsson B, Hedman E, *et al.* Predictors and moderators of Internet-based cognitive behavior therapy for obsessive-compulsive disorder: results from a randomized trial. *J Obsessive Compuls Relat Disord* 2015;4:1–7.
- 45 Andersson G. Prior treatments in a group of tinnitus sufferers seeking treatment. *Psychother Psychosom* 1997;66:107–10.
- 46 Fuhr K, Schröder J, Berger T, *et al.* The association between adherence and outcome in an Internet intervention for depression. *J Affect Disord* 2018;229:443–9.

## Supplementary File 1

Table 1: TRIPOD Checklist – Prediction model development

Section/Topic	Item	Checklist Item	Page
<b>Title and abstract</b>			
Title	1	Identify the study as developing and/or validating a multivariable prediction model, the target population, and the outcome to be predicted.	1
Abstract	2	Provide a summary of objectives, study design, setting, participants, sample size, predictors, outcome, statistical analysis, results, and conclusions.	2
<b>Introduction</b>			
Background and objectives	3a	Explain the medical context (including whether diagnostic or prognostic) and rationale for developing or validating the multivariable prediction model, including references to existing models.	3-5
	3b	Specify the objectives, including whether the study describes the development or validation of the model or both.	5
<b>Methods</b>			
Source of data	4a	Describe the study design or source of data (e.g., randomized trial, cohort, or registry data), separately for the development and validation data sets, if applicable.	6
	4b	Specify the key study dates, including start of accrual; end of accrual; and, if applicable, end of follow-up.	6
Participants	5a	Specify key elements of the study setting (e.g., primary care, secondary care, general population) including number and location of centres.	6
	5b	Describe eligibility criteria for participants.	6
	5c	Give details of treatments received, if relevant.	6-7
Outcome	6a	Clearly define the outcome that is predicted by the prediction model, including how and when assessed.	8-9
	6b	Report any actions to blind assessment of the outcome to be predicted.	NA
Predictors	7a	Clearly define all predictors used in developing or validating the multivariable prediction model, including how and when they were measured.	8-9
	7b	Report any actions to blind assessment of predictors for the outcome and other predictors.	NA
Sample size	8	Explain how the study size was arrived at.	9
Missing data	9	Describe how missing data were handled (e.g., complete-case analysis, single imputation, multiple imputation) with details of any imputation method.	10
Statistical analysis methods	10a	Describe how predictors were handled in the analyses.	10
	10b	Specify type of model, all model-building procedures (including any predictor selection), and method for internal validation.	10
	10d	Specify all measures used to assess model performance and, if relevant, to compare multiple models.	10
Risk groups	11	Provide details on how risk groups were created, if done.	NA
<b>Results</b>			
Participants	13a	Describe the flow of participants through the study, including the number of participants with and without the outcome and, if applicable, a summary of the follow-up time. A diagram may be helpful.	11
	13b	Describe the characteristics of the participants (basic demographics, clinical features, available predictors), including the number of participants with missing data for predictors and outcome.	11
Model development	14a	Specify the number of participants and outcome events in each analysis.	11
	14b	If done, report the unadjusted association between each candidate predictor and outcome.	11-12
Model specification	15a	Present the full prediction model to allow predictions for individuals (i.e., all regression coefficients, and model intercept or baseline survival at a given time point).	12-15
	15b	Explain how to use the prediction model.	14-15
Model performance	16	Report performance measures (with CIs) for the prediction model.	14-15
<b>Discussion</b>			
Limitations	18	Discuss any limitations of the study (such as nonrepresentative sample, few events per predictor, missing data).	18-19
Interpretation	19b	Give an overall interpretation of the results, considering objectives, limitations, and results from similar studies, and other relevant evidence.	15-18
Implications	20	Discuss the potential clinical use of the model and implications for future research.	19
<b>Other information</b>			
Supplementary information	21	Provide information about the availability of supplementary resources, such as study protocol, Web calculator, and data sets.	9, 11
Funding	22	Give the source of funding and the role of the funders for the present study.	19

**Supplementary File 2: Predictor Variables**

Table 1: Demographic variables

<b>Variable</b>	<b>Question</b>	<b>Response options</b>
Age	What is your age?	In years  Split into dichotomous variables (<=57 years of age and >57 years of age) based on the median
Gender	What is your gender?	Male (1), Female (2)
Education level	What is the highest level of education you have completed?	Highschool or less (1), College (2), Vocational training (3), Bachelor's degree (4), Master's degree or above (5)
Employment type	What best describes your employment?	Manager (1), Professional (2), Technical (3), Administrative (4), Skilled tradesman (5), Service occupation (6), Medical (7), Sales (8), Home maker (9), Student (10), Retired (11), Unemployed (12)
Loud noise exposure	Have you been exposed to loud noise?	Yes (1) , No (0)
Diagnosed with psychological condition	Have you been presently diagnosed with any psychological conditions including anxiety and depression?	Yes (1) , No (0)
Work less due to tinnitus	Do you work less because of your tinnitus?	No (0), Reduced hours (1), Stopped work (2), Disability allowance (3)

Table 2: Tinnitus and hearing-related variables

<b>Variable</b>	<b>Question</b>	<b>Response options</b>
Baseline tinnitus severity	Measured using the Tinnitus Functional Index (TFI)	Scores range from 0 to 100.  Split into dichotomous variables (<=55.2 and >55.2) based on the median
Tinnitus duration	How long have you had tinnitus for?	In years  Split into dichotomous variables (<=10.00 years and >10.00 years) based on the median
How often is tinnitus heard?	How often is tinnitus heard?	Occasionally (1), When taking out my hearing aid(s) (2), At night (3), Most of the time (4), All the time (5)
Tinnitus location	Where do you notice your tinnitus?	One ear (1), Both ears (2), In my head (3), Unsure (4), Other (5)

Type of tinnitus (9 different types)	<ul style="list-style-type: none"> <li>▪ Ringing</li> <li>▪ Buzzing</li> <li>▪ High pitched sound</li> <li>▪ Low pitched sound</li> <li>▪ Pulsing</li> <li>▪ Clicking</li> <li>▪ Music</li> <li>▪ Voices</li> <li>▪ Humming</li> </ul>	For each item: Yes (1) , No (0)
Multiple tones heard	This variable is computed based on responses to types of tinnitus. Answer yes to multiple types of tinnitus was considered as multiple tones heard	Yes (1) , No (0)
Presence of a hearing loss	Do you have a hearing loss?	No (0), Both ears (1), One ear (2), Unsure (3)

Table 3: Treatment-related variables

Variable	Question	Response options
Past treatment sought	Have you received treatment for tinnitus in the past?	Yes (1) , No (0)
Sounds can distract from tinnitus	How well can sounds around you distract you from your tinnitus or make the tinnitus less noticeable?	Fully (1), Partially (2), Not at all (3)
Hearing aid use	Do you wear hearing aid(s) or any other amplification devices?	No (0), One ear (1), Both ears (2)
Medication use	Do you currently take any medications?	Yes (1) , No (0)

Table 4: Clinical factors

Variable	Questionnaire	Number of items/ Response options	Score
Anxiety	General Anxiety Disorders (GAD-7)	7-items 4-point scale with “not at all” (score of 0) to “nearly every day” (score of 3)	Higher number indicates more severe anxiety (scores range between 0–21). The total score is interpreted as follows: <ul style="list-style-type: none"> <li>▪ 0–4: minimal anxiety</li> <li>▪ 5–9: mild anxiety</li> <li>▪ 10–14: moderate anxiety</li> <li>▪ 15–21: severe anxiety</li> </ul> Split into dichotomous variables ( $\leq 9$ no anxiety and $>9$ anxiety)
Depression	Patient Health Questionnaire (PHQ-9)	9-items 4-point scale with “not at all” (score of	Higher number indicates more severe depression (scores range between 0–27).

		0) to “nearly every day” (score of 3)	<p>The total score is interpreted as follows:</p> <ul style="list-style-type: none"> <li>▪ 5–9: mild depression</li> <li>▪ 10–14: moderate</li> <li>▪ 15–19: moderately severe</li> <li>▪ 20–18: severe depression</li> </ul> <p>Split into dichotomous variables (<math>\leq 14</math> no depression and <math>&gt; 14</math> depression)</p>
Insomnia	Insomnia Severity Index (ISA)	<p>7-item</p> <p>5-point scale with “no problem” (score of 0) to “very severe problem” (score of 4)</p>	<p>Higher number indicates more severe insomnia (scores range between 0–28).</p> <p>The total score is interpreted as follows:</p> <ul style="list-style-type: none"> <li>▪ 0–7: not clinically significant</li> <li>▪ 8–14: subthreshold insomnia</li> <li>▪ 15–21: clinical insomnia (moderate severity)</li> <li>▪ 22–28: clinical insomnia (severe degree)</li> </ul> <p>Split into dichotomous variables (<math>\leq 14</math> no insomnia and <math>&gt; 15</math> insomnia)</p>
Hyperacusis	Hyperacusis Questionnaire (HQ)	<p>14-items</p> <p>4-point scale with “no” (score of 0) to “yes, a lot” (score of 3)</p>	<p>Higher number more severe hyperacusis (scores range between 0–42).</p> <p>The total score is interpreted as follows:</p> <ul style="list-style-type: none"> <li>▪ <math>&gt; 28</math>: strong hypersensitivity</li> </ul> <p>Split into dichotomous variables (<math>\leq 28</math> no hyperacusis and <math>&gt; 28</math> hyperacusis)</p>
Hearing disability	Hearing Handicap Inventory for Adults – Screening (HHIA-S)	<p>10-items</p> <p>3-point scale with “yes” (score of 4) to “no” day (0)</p>	<p>Higher number more severe hearing disability (scores range between 0–40).</p> <p>The total score is interpreted as follows:</p> <ul style="list-style-type: none"> <li>▪ 0–8: no hearing disability</li> <li>▪ 10–24: mild to moderate hearing disability</li> <li>▪ 26–40: severe hearing disability</li> </ul> <p>Split into dichotomous variables (<math>\leq 8</math> no hearing disability and <math>\geq 10</math> hearing disability)</p>

Cognitive failures	Cognitive Failures Questionnaire (CFQ)	25-items  5-point scale with “never” (score of 0) to “very often” (score of 4)	Higher scores indicate more difficulties (cognitive failures) in perception, memory, and motor function (score range 0–100).  The total score is interpreted as follows: The scores range 0–100 with higher scores indicating more cognitive failures/problems (or reduced cognitive functioning).  Split into dichotomous variables (<=32 no cognitive problems and >32 cognitive problems)
Life satisfaction	Satisfaction with Life Scale (SWLS)	5-items  7-point scale with “strongly disagree” (score of 1) to “strongly agree” (7)	Higher number indicated more satisfaction with life (scores range between 5–35).  The total score is interpreted as follows: <ul style="list-style-type: none"> <li>▪ 0–9: extremely dissatisfied</li> <li>▪ 10–14: dissatisfied</li> <li>▪ 15–19: below average satisfaction</li> <li>▪ 20–24: average satisfaction</li> <li>▪ 25–29: high satisfaction</li> <li>▪ 30–35: highly satisfied</li> </ul> Split into dichotomous variables (<=19 life satisfaction and >19 high satisfaction)

### Supplementary File 3: Univariate analysis to examine association between predictor variables and outcome variable

Table 1: Participant demographic characteristics (n=228)

Characteristic	N (%)	Mean (SD)
<b>Demographic characteristics</b>		
Age (in years)		55.14 (12.92)
Gender		
▪ Female	98 (43%)	
▪ Male	130 (57%)	
Highest level of education		
▪ High school or below	59 (26%)	
▪ College	47 (21%)	
▪ Vocational training	31 (14%)	
▪ Bachelor's degree	61 (27%)	
▪ Masters degree or above	30 (13%)	
Employment		
▪ Manager	27 (12%)	
▪ Professional	46 (20%)	
▪ Technical	13 (6%)	
▪ Administrative	17 (7%)	
▪ Skilled tradesman	11 (5%)	
▪ Service occupation	11 (5%)	
▪ Medical	6 (3%)	
▪ Sales	8 (3%)	
▪ Homemaker	4 (2%)	
▪ Student	1 (0%)	
▪ Retired	73 (32%)	
▪ Unemployed	11 (5%)	
Loud noise exposure		
▪ Yes	103 (45%)	
▪ No	125 (55%)	
Diagnosed with a psychological condition		
▪ Yes	50 (22%)	
▪ No	178 (78%)	
Working less due to tinnitus		
▪ Reduced hours	8 (4%)	
▪ Stopped work	32 (14%)	
▪ Disability allowance	7 (3%)	
▪ No	181 (79%)	
<b>Tinnitus and hearing-related characteristics</b>		
Baseline tinnitus severity (measured using Tinnitus Functional Index)		57.93 (19.17)
Tinnitus duration (in years)		17.68 (19.42)
How often tinnitus is heard		

<ul style="list-style-type: none"> <li>▪ Occasionally</li> <li>▪ When taking out my hearing aid(s)</li> <li>▪ At night</li> <li>▪ Most of the time</li> <li>▪ All the time</li> </ul>	4 (2%) 3 (1%) 4 (2%) 63 (27%) 154 (68%)	
Tinnitus location <ul style="list-style-type: none"> <li>▪ One ear</li> <li>▪ Both ears</li> <li>▪ In my head</li> <li>▪ Other location</li> <li>▪ Unsure</li> </ul>	61 (27%) 109 (48%) 34 (15%) 3 (1%) 21 (9%)	
Type of tinnitus sound (answering Yes) <ul style="list-style-type: none"> <li>▪ Ringing</li> <li>▪ Buzzing</li> <li>▪ High pitched sound</li> <li>▪ Low pitched sound</li> <li>▪ Pulsating</li> <li>▪ Clicking</li> <li>▪ Music</li> <li>▪ Voices</li> <li>▪ Humming</li> </ul>	71 (31%) 75 (33%) 130 (57%) 16 (7%) 28 (12%) 14 (6%) 4 (2%) 3 (1%) 21 (9%)	
Multiple sounds heard <ul style="list-style-type: none"> <li>▪ Yes</li> <li>▪ No</li> </ul>	73 (32%) 155 (68%)	
Presence of a hearing loss <ul style="list-style-type: none"> <li>▪ No</li> <li>▪ Both ears</li> <li>▪ One ear</li> <li>▪ Unsure</li> </ul>	49 (21%) 104 (46%) 46 (20%) 29 (13%)	
<b>Treatment-related characteristics</b>		
Past tinnitus treatment sought <ul style="list-style-type: none"> <li>▪ Yes</li> <li>▪ No</li> </ul>	58 (25%) 170 (75%)	
Sounds can distract from tinnitus <ul style="list-style-type: none"> <li>▪ Fully</li> <li>▪ Partially</li> <li>▪ Not at all</li> </ul>	26 (11%) 178 (78%) 24 (10%)	
Hearing aid use <ul style="list-style-type: none"> <li>▪ No</li> <li>▪ Unilateral</li> <li>▪ Bilateral</li> </ul>	159 (70%) 19 (8%) 50 (22%)	
Medication use <ul style="list-style-type: none"> <li>▪ Yes</li> <li>▪ No</li> </ul>	130 (57%) 98 (43%)	



Table 2: Univariate analysis with the Chi-square/ Fishers exact test results on the association between the demographic predictor categories and outcome variable (success as defined by a TFI-score changes  $\geq 13$  points or a failure). \*Indicates use of Fisher's exact test results due to less than 5 cases in subcategories.

Predictor variable	Sub-Categories	Crude Odds Ratio (95% CIs)	P-Value
Age	>57 years	0.85 (0.50, 1.47)	0.57
	$\leq 57$ years	Ref	
Gender	Female	1.12 (0.64, 1.94)	0.70
	Male	Ref	
Education level	College	0.61 (0.31, 1.42)	<b>0.01*</b>
	Vocational training	1.70 (0.75, 4.88)	
	Bachelor's degree	1.30 (0.67, 2.92)	
	<b>Master's degree or above</b>	<b>3.47 (1.32, 12.51)</b>	
	High school or less	Ref	
Employment type	Professional	0.59 (0.25, 1.82)	0.95*
	Technical	0.40 (0.13, 1.89)	
	Administrative	0.40 (0.14, 1.66)	
	Skilled tradesman	0.56 (0.18, 3.00)	
	Service occupation	0.80 (0.24, 4.66)	
	Medical	1.00 (0.22, 11.54)	
	Sales	0.80 (0.21, 6.00)	
	Home maker	0.27 (0.06, 3.00)	
	Student	0.40 (0.05, 35.47)	
	Retired	0.74 (0.32, 2.12)	
	Unemployed	0.80 (0.24, 4.66)	
	Manager	Ref	
Loud noise exposure	Yes	0.80 (0.46, 1.38)	0.43
	No	Ref	
Presence of a psychological condition	Yes	1.72 (0.85, 3.46)	0.13
	No	Ref	
Work less due to tinnitus	Reduced hours	1.05 (0.31, 6.18)	0.89*
	Stopped work	0.81 (0.41, 1.89)	
	Disability allowance	0.53 (0.16, 2.88)	
	No	Ref	

Table 3: Univariate analysis with the Chi-square/ Fishers exact test results on the association between the tinnitus and hearing-related predictor categories and outcome variable (success as defined by a TFI-score changes  $\geq 13$  points or a failure). \*Indicates use of Fisher's exact test results due to less than 5 cases in subcategories.

Predictor variable	Sub-Categories	Crude Odds Ratio (95% CIs)	P-Value
Baseline tinnitus severity	>55.2	<b>2.65 (1.50, 4.67)</b>	<b>0.001</b>
	$\leq 55.2$	Ref	
Tinnitus duration	>10.00 years	1.16 (0.66, 2.02)	0.60
	$\leq 10.00$ years	Ref	
How often tinnitus is heard	When taking out my hearing aid(s)	0.67 (0.02, 18.06)	0.19*
	At night	0.33 (0.02, 6.65)	
	Most of the time	0.39 (0.04, 3.96)	
	All the time	0.76 (0.08, 7.49)	
	Occasionally	Ref	
Tinnitus location	Both ears	1.41(0.48, 4.16)	0.90*
	In my head	0.94 (0.48, 1.80)	
	Unsure	1.35 (0.55, 3.34)	
	Other	1.13 (0.10,13.16)	
	One ear	Ref	
Tinnitus type: Ringing	Yes	1.30 (0.72, 2.37)	0.38
	No	Ref	
Tinnitus type: Buzzing	Yes	1.34 (0.74, 2.42)	0.32
	No	Ref	
Tinnitus type: High pitch	Yes	0.76 (0.44, 1.33)	0.34
	No	Ref	
Tinnitus type: Low pitch	Yes	0.89 (0.31, 2.56)	0.83
	No	Ref	
Tinnitus type: Pulsing	Yes	0.97 (0.42, 2.21)	0.94
	No	Ref	
Tinnitus type: Clicking	Yes	0.52 (0.17, 1.53)	0.23
	No	Ref	
Tinnitus type: Music	Yes	1.63 (0.17, 15.98)	1.00*
	No	Ref	
Tinnitus type: Voices	Yes	0.09 (0.00, 1.75)	0.04*
	No	Ref	
Tinnitus type: Humming	Yes	0.56 (0.23, 1.39)	0.21
	No	Ref	
Multiple tones heard	Yes	1.15 (0.64, 2.08)	0.63
	No	Ref	
Presence of a hearing loss	Both ears	1.20 (0.59, 2.41)	0.92
	One ear	1.19 (0.51, 2.74)	
	Unsure	1.41 (0.53, 3.73)	
	No	Ref	

Table 4: Univariate analysis with the Chi-square/ Fishers exact test results on the association between the treatment-related predictor categories and outcome variable (success as defined by a TFI-score changes  $\geq 13$  points or a failure). \*Indicates use of Fisher's exact test results due to less than 5 cases in subcategories.

Predictor variable	Sub-Categories	Crude Odds Ratio (95% CIs)	P-Value
Past treatment sought	Yes	0.94 (0.50, 1.74)	0.83
	No	Ref	
Sounds can distract	Partially	<b>4.34 (1.82, 10.34)</b>	<b>0.001</b>
	Not at all	<b>3.15 (0.99, 10.00)</b>	
	Fully	Ref	
Hearing aid use	One ear	<b>1.57 (0.61, 5.49)</b>	<b>0.26</b>
	Both ear	0.69 (0.38, 1.39)	
	No	Ref	
Medication use	Yes	1.22 (0.71, 2.12)	0.46
	No	Ref	

Table 5: Univariate analysis with the Chi-square/ Fishers exact test results on the association between the clinical factors predictor categories and outcome variable (success as defined by a TFI-score changes  $\geq 13$  points or a failure). \*Indicates use of Fisher's exact test results due to less than 5 cases in subcategories.

Predictor variable	Sub-Categories	Odds Ratio (95% CIs)	P-Value
Anxiety	Yes	1.53 (0.83, 2.82)	0.17
	No	Ref	
Depression	Yes	1.54 (0.62, 3.83)	0.35
	No	Ref	
Insomnia	Yes	1.27 (0.72, 2.23)	0.41
	No	Ref	
Hyperacusis	Yes	1.21 (0.56, 2.63)	0.62
	No	Ref	
Hearing disability	Yes	1.37 (0.77, 2.43)	0.28
	No	Ref	
Cognitive functions	Yes	0.99 (0.56, 1.74)	0.97
	No	Ref	
Life satisfaction	Yes	0.76 (0.44, 1.33)	0.34
	No	Ref	

**Supplementary File 1**

Table 1: TRIPOD Checklist – Prediction model development

Section/Topic	Item	Checklist Item	Page
<b>Title and abstract</b>			
Title	1	Identify the study as developing and/or validating a multivariable prediction model, the target population, and the outcome to be predicted.	1
Abstract	2	Provide a summary of objectives, study design, setting, participants, sample size, predictors, outcome, statistical analysis, results, and conclusions.	1
<b>Introduction</b>			
Background and objectives	3a	Explain the medical context (including whether diagnostic or prognostic) and rationale for developing or validating the multivariable prediction model, including references to existing models.	1
	3b	Specify the objectives, including whether the study describes the development or validation of the model or both.	1
<b>Methods</b>			
Source of data	4a	Describe the study design or source of data (e.g., randomized trial, cohort, or registry data), separately for the development and validation data sets, if applicable.	2
	4b	Specify the key study dates, including start of accrual; end of accrual; and, if applicable, end of follow-up.	2
Participants	5a	Specify key elements of the study setting (e.g., primary care, secondary care, general population) including number and location of centres.	2
	5b	Describe eligibility criteria for participants.	2
	5c	Give details of treatments received, if relevant.	2-3
Outcome	6a	Clearly define the outcome that is predicted by the prediction model, including how and when assessed.	3
	6b	Report any actions to blind assessment of the outcome to be predicted.	NA
Predictors	7a	Clearly define all predictors used in developing or validating the multivariable prediction model, including how and when they were measured.	3
	7b	Report any actions to blind assessment of predictors for the outcome and other predictors.	NA
Sample size	8	Explain how the study size was arrived at.	2
Missing data	9	Describe how missing data were handled (e.g., complete-case analysis, single imputation, multiple imputation) with details of any imputation method.	3
Statistical analysis methods	10a	Describe how predictors were handled in the analyses.	3
	10b	Specify type of model, all model-building procedures (including any predictor selection), and method for internal validation.	3
	10d	Specify all measures used to assess model performance and, if relevant, to compare multiple models.	3
Risk groups	11	Provide details on how risk groups were created, if done.	NA
<b>Results</b>			
Participants	13a	Describe the flow of participants through the study, including the number of participants with and without the outcome and, if applicable, a summary of the follow-up time. A diagram may be helpful.	3
	13b	Describe the characteristics of the participants (basic demographics, clinical features, available predictors), including the number of participants with missing data for predictors and outcome.	3
Model development	14a	Specify the number of participants and outcome events in each analysis.	3
	14b	If done, report the unadjusted association between each candidate predictor and outcome.	3
Model specification	15a	Present the full prediction model to allow predictions for individuals (i.e., all regression coefficients, and model intercept or baseline survival at a given time point).	3-5
	15b	Explain how to use the prediction model.	3-5
Model performance	16	Report performance measures (with CIs) for the prediction model.	3-5
<b>Discussion</b>			
Limitations	18	Discuss any limitations of the study (such as nonrepresentative sample, few events per predictor, missing data).	7
Interpretation	19b	Give an overall interpretation of the results, considering objectives, limitations, and results from similar studies, and other relevant evidence.	6-7
Implications	20	Discuss the potential clinical use of the model and implications for future research.	7
<b>Other information</b>			
Supplementary information	21	Provide information about the availability of supplementary resources, such as study protocol, Web calculator, and data sets.	2
Funding	22	Give the source of funding and the role of the funders for the present study.	8